



INTEGRAL TAU METHOD FOR FOURTH-ORDER INITIAL VALUE PROBLEMS WITH THIRD DEGREES OF OVER - DETERMINATION

Ojo Victoria Oluwatoyin¹, Adeniyi Raphael Babatunde² and Adeyefa Emmanuel³

¹Department of Statistics, Oyo State College of Agriculture and Technology, Igbo-Ora, Oyo State, Nigeria.

²Department of Mathematics, University of Ilorin, Ilorin, Nigeria.

³ Department of Mathematics, Federal University, Oye Ekiti, Ekiti State, Nigeria.

ABSTRACT: This paper is concerned with the solution of a class of fourth order initial value problems in ordinary differential equation by the integrated formulation of the tau method. The initial focus is the class with a maximum of third degree overdetermination. The matrix equations were constructed based on the degree of overdetermination and for purpose of automation. The automated Tau System was tested on some selected problems to validate the study numerical evidences, thus obtained, confirm the accuracy of the method.

KEYWORDS: Tau Method, Variant, Formulation, Approximant, Perturbation Term

INTRODUCTION

Lanczos proposed the tau method techniques in 1983 for the numerical solution of ordinary differential equation with some conditions given as

$$Ly_n(x) = \sum_{r=0}^m \left(\sum_{k=0}^N P_{rk} x^k \right) y^{(r)}(x) = \sum_{r=0}^n f_r x^r \quad a \leq x \leq b \quad (1.1)$$

$$L * Y(x_{rk}) = \sum_{r=0}^{m-1} a_{rk} y^{(r)}(x_{rk}) = \alpha_k \quad k = 1(1)m \quad (1.2)$$

By seeking an approximate solution of the form:

$$Y_n(x) = \sum_{r=0}^n a_r x^r \quad (1.3)$$

$r < +\infty$ of $y(x)$ which is the exact solution of the corresponding perturbed system

$$L * Y_n(x) = \sum_{r=0}^n f_r x^r H_n(x) \quad (1.4)$$

$$LY_n(x_{nk}) = \alpha_k \quad k = 1(1)m \quad (1.5)$$



Where

L is the linear differential operator $\alpha_k, f_r, P_{nk}, -N_M : r = 0 \text{ to } m, K = 0 \text{ to } N_r$, a and b are real constants, $y(r)$ denotes the derivatives of order r of $y(x)$. The perturbation term $H_n(x)$ in 1.4 is defined by

$$H_n(x) = \sum_{i=0}^{m+s-1} \tau_{i+1} T_{n-m+i+1}(x) = \sum_{i=0}^{m+s-1} \tau_{i+1} \sum_{r=0}^{n-m+i+1} C_r^{(n-m+i+1)} x^r \quad (1.6)$$

And $C_r^{(n)}$'s are the coefficient of power of x (that is x^r) in the n th degree chebyshev polynomial denoted and defined by

$$T_n(x) = \cos \left(n \cos^{-1} \left[\frac{2x - a - b}{b - a} \right] \right) = \sum_{r=0}^n C_r^{(n)} \quad (1.7)$$

The r 's are the free tau parameters to be determined alongside with a_r and S is the number of over-determinations of (1.1), which is defined by

$$S = \max[N_r - r : 0 \leq r \leq m, N_r \geq r] \geq 0 \quad (1.8)$$

LITERATURE REVIEW

The Tau method was initially formulated as a tool for the approximation of special function of mathematical physics which could be expressed in terms of simple differential equations. It later developed into a powerful and accurate tool for the numerical solution of complex differential and functional equations. The main idea in it is to solve approximate problem. Accurate approximate polynomial solution in a linear ordinary differential equation with polynomial coefficient can be obtained by the Tau method introduced in [1]. The method is related to the principle of economization of a differentiable function implicitly defined by a linear differential equation with polynomial coefficient. Techniques based on the Tau method have been reported in the literature with application to more general equations including non-linear ones [2-3], while techniques based on direct Chebyshev replacement have been discussed in [4] and more recently in the work of [5]. Further details on the Tau method can be found in reference [6-13]. Because of the limitation in some of the works [7], this study seeks to extend the scope to fourth order problems with third degree overdetermination.

The Integrated Formulation of the TAU Method

Description of the integrated formulation

Let us consider the m -th order linear differential system

$$Ly(x) := \sum_{r=0}^m \alpha_r(x) y^{(r)}(x) = \sum_{r=0}^f f_r x^r, a \leq x \leq b \quad (9)$$



$$L^*y(x_{rk}) := \sum_{r=0}^{m-1} \alpha_{rk} y^{(r)}(x_{rk}) = \alpha_k, \quad k = 1(1)m \quad (2.2)$$

Let $\int \int \int \cdots \int^i y(x) dx$ denote the indefinite integration i times applied to the function $g(x)$ and let

$$I_L = \int \int \int \cdots \int^m L(\cdot) dx \quad (2.3)$$

The integral form of (2.3) now becomes

$$I_L(y(x)) = \int \int \int \cdots \int^m f(x) dx + C_{m-1}(x) \quad (2.4)$$

The tau approximant $y_n(x)$ of (2.1), satisfies the perturbed problem:

$$I_L(y_n(x)) = \int \int \int \cdots \int^m f(x) dx + H_{n+m+1}(x) \quad (2.5)$$

$$L^*y_n(x_{rk}) = \alpha_k, k = 1(1)m \quad (2.6)$$

where:

$$H_{n+m}(x) = \sum_{r=0}^{m+s+1} \tau_{m+s-r} T_{n-m+r+1}(x) \quad (2.7)$$

A Class of Overdetermined Fourth Order Differential Equations

We consider here the integrated form of the tau method for the class of problems:

$$Ly(x) := \sum_{r=0}^m P_r(x) y^{(r)}(x) = f(x), \quad a \leq x \leq b \quad (3.1.0)$$

$$L^*y_n(x_{rk}) = \sum_{r=0}^{m-1} a_{rk} y_n^{(r)}(x_{rk}) = \alpha_k, \quad k = 0(1)(m-1) \quad (3.1.2)$$

$$P_r(x) = \sum_{k=0}^{N_r} p_{rk} x^k \quad (3.1.3)$$

for the case $m = 4$ and $s = 3$.

So, we now derive a fifth degree approximants for the equation. From (3.1.0), the general case for $m = 4$, $s = 3$ is given by



$$\begin{aligned}
 Ly(x) &:= (\alpha_0 + \alpha_1 x + \alpha_2 x^2 + \alpha_3 x^3 + \alpha_4 x^4 + \alpha_5 x^5 + \alpha_6 x^6 + \alpha_7 x^7) y^{iv}(x) + \beta_0 + \beta_1 x \\
 &\quad + \beta_2 x^2 + \beta_3 x^3 + \beta_4 x^4 + \beta_5 x^5 + \beta_6) y^{iii}(x) \\
 &\quad + (\gamma_0 + \gamma_1 x + \gamma_2 x^2 + \gamma_3 x^3 + \gamma_4 x^4 + \gamma_5 x^5) y^{ii}(x) (\lambda_0 + \lambda_1 x + \lambda_2 x^2 + \lambda_3 x^3 \\
 &\quad + \lambda_4 x^4) y^i(x) \\
 &\quad + (\mu_0 + \mu_1 x + \mu_2 x^2 \\
 &\quad + \mu_3 x^3) y(x) = f(x) = \sum_{r=0}^n f_r x^r
 \end{aligned} \tag{3.1.4}$$

$$y(0) = \rho_0, y'(0) = \rho_1, y''(0) = \rho_2, y'''(0) = \rho_3$$

where, for convenience, we have chosen $\alpha, \beta, \gamma, \lambda$ and μ to denote $\rho_4, \rho_3, \rho_2, \rho_1$ and ρ_0 respectively; $x_{rk} = 0$ and $a = 0$. That is,

$$\begin{aligned}
 Ly(x) &:= \int_0^x \int_0^u \int_0^t \int_0^s (\alpha_0 + \alpha_1 v + \alpha_2 v^2 + \alpha_3 v^3 + \alpha_4 v^4 + \alpha_5 v^5 + \alpha_6 v^6 + \alpha_7 v^7) y^{iv}(v) dv ds dt du + \int_0^x \int_0^u \int_0^t \int_0^s \\
 &\quad (\beta_0 + \beta_1 v + \beta_2 v^2 + \beta_3 v^3 + \beta_4 v^4 + \beta_5 v^5 + \beta_6) y'''(v) dv ds dt du + \int_0^x \int_0^u \int_0^t \int_0^s (\gamma_0 + \gamma_1 v + \gamma_2 v^2 + \gamma_3 v^3 + \\
 &\quad \gamma_4 v^4 + \gamma_5 v^5) y''(v) dv ds dt du + \int_0^x \int_0^u \int_0^t \int_0^s (\lambda_0 + \lambda_1 v + \lambda_2 v^2 + \lambda_3 v^3 + \lambda_4 v^4) y'(v) dv ds dt du + \\
 &\quad \int_0^x \int_0^u \int_0^t \int_0^s (\mu_0 + \mu_1 v + \mu_2 v^2 + \mu_3 v^3) y(v) dv ds dt du = \int_0^x \int_0^u \int_0^t \int_0^s f(V) dv ds dt du + \tau_1 T_{n+7}(x) \\
 &\quad + \tau_1 T_{n+6}(x) + \tau_2 T_{n+5}(x) + \tau_3 T_{n+4}(x) + \tau_4 T_{n+3}(x) + \tau_5 T_{n+2}(x) + \tau_6 T_{n+1}(x)
 \end{aligned} \tag{3.1.5}$$

After simplifying and equating the Corresponding Coefficient Powers of x, we have the recurrence relation:

$$\alpha_0 a_0 - \tau_1 C_0^{n+7} - \tau_2 C_0^{n+6} - \tau_3 C_0^{n+5} - \tau_4 C_0^{n+4} - \tau_5 C_0^{n+3} - \tau_6 C_0^{n+2} - \tau_7 C_0^{n+1} = \alpha_0 \rho_0 \tag{3.1.6}$$

$$\beta_0 a_0 + \alpha_0 a_1 - \tau_1 C_1^{n+7} - \tau_2 C_1^{n+6} - \tau_3 C_1^{n+5} - \tau_4 C_1^{n+4} - \tau_5 C_1^{n+3} - \tau_6 C_1^{n+2} - \tau_7 C_1^{n+1} = \alpha_1 \rho_1 - 2\alpha_0 \rho_1 - 2\beta_0 \rho_0 \tag{3.1.7}$$

$$\begin{aligned}
 \frac{1}{2}[(3\gamma_0 - 4\beta_1 - 7\alpha_2)a_0 - (\alpha_1 + \beta_0)a_1 - 2\alpha_0 a_0] - \tau_1 C_2^{n+7} - \tau_2 C_2^{n+6} - \tau_3 C_2^{n+5} - \tau_4 C_2^{n+4} - \tau_5 C_2^{n+3} - \tau_6 C_2^{n+2} - \tau_7 C_2^{n+1} \\
 = \frac{\alpha_0 \rho_1 + 2\alpha_2 \beta_2 - \beta_0 \gamma_0 - 3\gamma_0 \rho_0 - 4\beta_1 \rho_0 - 7\alpha_2 \rho_0 + \alpha_1 \rho_1 + 2\alpha_0 \rho_2}{2}
 \end{aligned} \tag{3.1.8}$$



$$\frac{1}{6}[(30\alpha_3 + 12\beta_2 + 5\gamma_1 + 2\lambda_0)a_0 + (8\alpha_2 + 5\beta_1 + 3\gamma_0)a_1 - 2(2\alpha_1 + \beta_0)a_2 + \alpha_0a_3] - \tau_1C_3^{n+7} - \tau_2C_3^{n+6}$$

$$-\tau_3C_3^{n+5} - \tau_4C_3^{n+4} - \tau_5C_3^{n+3} - \tau_6C_3^{n+2} - \tau_7C_3^{n+1}$$

$$= \frac{181\alpha_3\rho_0 - 3\alpha_1\rho_2 + 6\alpha_2\rho_1 - 3\beta_0\rho_2 + 10\beta_2\rho_0 + 6\gamma_1\rho_0 + 2\gamma_0\rho_1 + \lambda_0\rho_0 - 5\alpha_0\rho_3 + 3\rho_1 + 5\beta_1\rho_1}{6} \quad (3.1.9)$$

$$\frac{1}{24}[(116\alpha_4 + 30\beta_3 + 8\gamma_2 + \lambda_1 + \mu_0)a_0 + (42\alpha_3 + 14\beta_2 + 4\gamma_1 + 2\lambda_0)a_1 + (18\alpha_2 + 12\beta_1 + 6\gamma_0)a_2 - (18\alpha_1 + 6\beta_0)$$

$$a_3 + 24\alpha_0a_4] - \tau_1C_4^{n+7} - \tau_2C_4^{n+6} - \tau_3C_4^{n+5} - \tau_4C_4^{n+4} - \tau_5C_4^{n+3} - \tau_6C_4^{n+2} - \tau_7C_4^{n+1} = 0 \quad (3.2.0)$$

$$\frac{1}{120}[(720\alpha_5 + 168\beta_4 + 368\gamma_3 + 6\lambda_2 + \mu_1)a_0 + (264\alpha_4 + 96\beta_3 + 24\gamma_4 + 3\lambda_1 + \mu_0)a_1 + 2(42\alpha_3 + 34\beta_2 + 11\gamma_1$$

$$+ 2\lambda_0)a_2 + (60\alpha_2 + 42\beta_1 + 18\gamma_0)a_3 + 24(4\alpha_1 + \beta_0)a_4 + 120\alpha_0a_5] - \tau_1C_5^{n+7} - \tau_2C_5^{n+6} - \tau_3C_5^{n+5} - \tau_4C_5^{n+4} - \tau_5C_5^{n+3} - \tau_6C_5^{n+2} - \tau_7C_5^{n+1} = 0 \quad (3.2.1)$$

$$\frac{1}{360}[(1440\alpha_6 + 900\beta_5 + 48\gamma_4 + 24\lambda_3 + \mu_2)a_0 + (1240\alpha_5 + 300\beta_4 + 60\gamma_3 + 8\lambda_2 + \mu_1)a_1 + (432\alpha_1 - 198\beta_3 -$$

$$46\gamma_2 - 5\lambda_1 + \mu_0)a_2 + (72\alpha_3 + 144\beta_2 + 42\gamma_1 + 6\lambda_0)a_3 + 120(11\alpha_2 - 8\beta_1 - 3\gamma_0)a_4 + 60(5\alpha_1 + \beta_0)a_5 +$$

$$360\alpha_0a_6] - \tau_1C_6^{n+7} - \tau_2C_6^{n+6} - \tau_3C_6^{n+5} - \tau_4C_6^{n+4} - \tau_5C_6^{n+3} - \tau_6C_6^{n+2} - \tau_7C_6^{n+1} = 0 \quad (3.2.2)$$

$$[\frac{[\alpha_1k - \alpha_1 + \beta_0]}{k}a_{k-1} + \frac{[\alpha_2(k-1) - (2\alpha_2 + \beta_1)(k-1) - 3(2\alpha_2 + \beta_1 + \gamma_0)]}{k(k-1)}a_{k-2} + [\alpha_3(k-2)$$

$$\frac{(k-1)k - (3\alpha_3 + \beta_2)(k-2)(k-1) - 3(6\alpha_3 + 2\beta_2 + \gamma_1)(k-2) - 2(6\alpha_3 + 2\beta_2 + \gamma_1 - \lambda_0)}{(k-2)(k-1)k}a_{k-3}$$

$$[\alpha_4(k-3)(k-2)(k-1)k - (4\alpha_4 + \beta_3)(k-3)(k-2)(k-1) - 3(12\alpha_4 + 3\beta_3 + \gamma_2)$$



$$\frac{(k-3)(k-2)-2(24\alpha_4+6\beta_3+2\gamma_2-\lambda_1)(k-3)+(24\alpha_4+6\beta_3+2\gamma_2+\lambda_1+\mu_0)]}{(k-3)(k-2)(k-1)k}a_{k-4}$$

$$[\alpha_5(k-3)(k-2)(k-1)k-(5\alpha_5+\beta_4)(k-3)(k-2)(k-1)-3(20\alpha_5+4\beta_4+\gamma_3)$$

$$\frac{(k-3)(k-2)-2(60\alpha_5+12\beta_4+3\gamma_3-\lambda_2)(k-3)-(120\alpha_5+24\beta_4+6\gamma_3+2\lambda_2+\mu_1)]}{(k-3)(k-2)(k-1)k}a_{k-5}$$

$$[\alpha_6(k-3)(k-2)(k-1)k-(6\alpha_6+\beta_5)(k-3)(k-2)(k-1)-3(30\alpha_6+5\beta_5+\gamma_4)$$

$$\frac{(k-3)(k-2)-2(120\alpha_6+120\beta_5+4\gamma_4+\lambda_3)(k-3)+(360\alpha_6+60\beta_5+12\gamma_4+3\lambda_3+\mu_2)]}{(k-3)(k-2)(k-1)k}a_{k-6}$$

$$[\alpha_7(k-3)(k-2)(k-1)k-(7\alpha_7+\beta_6)(k-3)(k-2)(k-1)-3(42\alpha_7+6\beta_6+\gamma_5)$$

$$\frac{(k-3)(k-2)-2(210\alpha_7+30\beta_6+5\gamma_5+\lambda_4)(k-3)+(840\alpha_7+120\beta_6+20\gamma_5+4\lambda_4+\mu_3)]}{(k-3)(k-2)(k-1)k}a_{k-7}$$

$$\begin{aligned} & \tau_1 C_K^{n+7} - \tau_2 C_K^{n+6} - \tau_3 C_K^{n+5} - \tau_4 C_K^{n+4} - \tau_5 C_K^{n+3} - \tau_6 C_K^{n+2} - \tau_7 C_K^{n+1} \\ &= \frac{f_{k-7}}{(k)(k-1)(k-2)(k-3)(k-4)(k-5)(k-6)} \end{aligned} \quad (3.2.3)$$

$$\frac{1}{(n-2)(n-1)(n)(n+1)}[[\alpha_7(n-2)[n^3-4n^2-120n-294]+840\alpha_7)-(\beta_6(n-2)[n^2+17n+62]-$$

$$120\beta_6)-(\gamma_5(n-2)(n-1))-20\gamma_5)-(2\lambda_4(n-4)+\mu_3)a_{n-6}]+[(\alpha_6(n-2)(n^3-6n^2-85n-150)+360\alpha_6)$$

$$-(\beta_5(n-2)[n^2+14n+225]-60\beta_5)-\gamma_4(n-2)[3n+5]-12\gamma_4)-2\lambda_3(n-17)+\mu_2)]a_{n-5}+[(\alpha_5(n-2)$$

$$[n^3-5n^2-56n-60]-120\alpha_5)-(\beta_4(n-2)(n^2+11n+12)-24\beta_4)-\gamma_3(n-2)(3n+3)+6\gamma_3)+\lambda_2n+\mu_1)]$$



$$a_{n-4} + [\alpha_4(n-2)[-35n^3 + 20n^2 + 15n] + 24\alpha_4] - [\beta_3(n-2)[n^2 - 10n - 3] + 6\beta_3] + [\gamma_2(n-2)(-3n-1)$$

$$+ 2\gamma_2] + [\lambda_1(5-2n) + \mu_0]a_{n-3} + [(\alpha_3(n-2)(n^3 - 3n^2 - 19n + 6)] + \beta_2[2n^2 + 10n] + \gamma_1(1-3n) - 2\lambda_0]$$

$$a_{n-2} + [(\alpha_2(n-2)(n^2 - 7n + 6)] - \beta_1[n+3] - 3\gamma_0]a_{n-1} + [\alpha_1(n-2)(n^2 - n)(n + \beta_0)]a_n]$$

$$- \tau_1 C_{n+1}^{n+7} - \tau_2 C_{n+1}^{n+6} - \tau_3 C_{n+1}^{n+5} - \tau_4 C_{n+1}^{n+4} - \tau_5 C_{n+1}^{n+3} - \tau_6 C_{n+1}^{n+2} - \tau_7 C_{n+1}^{n+1}]x^{n+1}$$

$$= \frac{f_{n-6}}{(n+1)(n)(n-1)(n-2)(n-3)(n-4)(n-5)} \quad (3.2.4)$$

$$\frac{1}{(n-1)(n)(n+1)(n+2)} [[(\alpha_7(n-1)[n^3 - 4n^2 - 131n - 420] + 840\alpha_7) - (\beta_6(n-1)[n^2 + 19n + 60] -$$

$$120\beta_6) - (\gamma_5(n-1)(3n+10)] - 20\gamma_5) - (2\lambda_4(n-3) + \mu_3)a_{n-5}] + [(\alpha_6(n-1)(n^3 - 3n^2 - 86n - 240) +$$

$$360\alpha_6) - (\beta_5(n-1)[n^2 + 16n + 240] - 60\beta_5) - \gamma_4(n-1)[3n+8] - 12\gamma_4) - 2\lambda_3(n+6) + \mu_2)]a_{n-4} + [(\alpha_5$$

$$(n-1)[n^3 - 2n^2 - 63n - 120] - 120\alpha_5) - (\beta_4(n-1)(n^2 + 13n + 24) + 24\beta_4) - \gamma_3(n-1)(3n+6) + 6\gamma_3) -$$

$$2\lambda_2n - \mu_1)]a_{n-3} + [\alpha_4(n+1)[n^3 - 39n^2 + 38n - 48] - 24\alpha_4] - (\beta_3(n-1)[n^2 - 2n] + 6\beta_3) - (\gamma_2(n-1)$$

$$(3n+4) - 2\gamma_2) + (\lambda_1(2n+3) + \mu_0)]a_{n-2} + [(\alpha_3(n-1)(n^3 - 19n) - 12\alpha_3) - \beta_2(n^2 - 7n) - 4\beta_2)$$

$$(\gamma_1(3n+2) - 2\lambda_0)]a_{n-1} + [(\alpha_2(n-1)(n^2 - 7n) - \beta_1[n+4] - 3\gamma_0]a_n - \tau_1 C_{n+2}^{n+7} - \tau_2 C_{n+2}^{n+6} - \tau_3 C_{n+2}^{n+5}$$

$$- \tau_4 C_{n+2}^{n+4} - \tau_5 C_{n+2}^{n+3} - \tau_6 C_{n+2}^{n+2}] = \frac{f_{n-5}}{(n+2)(n+1)(n)(n-1)(n-2)(n-3)(n-4)} \quad (3.2.5)$$



$$\begin{aligned}
& \frac{1}{(n)(n+1)(n+2)(n+3)} [[(\alpha_7 n[n^3 - n^2 - 136n - 554] + 840\alpha_7) - (\beta_6 n)[n^2 + 21n + 80] - \\
& 120\beta_6) - (\gamma_5 n(3n - 7)] - 20\gamma_5) - 2\lambda_4(n - 2) + \mu_3)] a_{n-4} + [(\alpha_6 n(n^3 - 85n - 336) + \\
& 360\alpha_6) - (\beta_5 n(n^2 + 18n + 25) - 60\beta_5) - (\gamma_4 n(3n - 5) - 12\gamma_4) - 2\lambda_3(n - 15) + \mu_2)] a_{n-3} + [(\alpha_5 n \\
& [n^3 + n^2 - 64n - 184] - 120\alpha_5) - (\beta_4 n(n^2 + 15n + 38) + 24\beta_4) - \gamma_3 n(3n + 9) + 6\gamma_3) - 2\lambda_2(n + 1) - \mu_1)] a_{n-2}
\end{aligned}$$

$$\begin{aligned}
& a_{n-2} + [\alpha_4 n[n^3 + 2n^2 - 37n - 86] + 24\alpha_4) - (\beta_3 n(n^2 + 12n + 23) - (\gamma_2 n(3n + 7) - 2\gamma_2) \\
& - (\lambda_1(2n - 1) + \mu_0)] a_{n-1} + [(\alpha_3 n(n^3 + 3n^2 - 16n - 30) - \beta_2 n(n + 9) - 12\beta_2) - (\gamma_1(3n + 5) - 2\lambda_0)] a_n \\
& - \tau_1 C_{n+3}^{n+7} - \tau_2 C_{n+3}^{n+6} - \tau_3 C_{n+3}^{n+5} - \tau_4 C_{n+3}^{n+4} - \tau_5 C_{n+3}^{n+3} \\
& = \frac{f_{n-4}}{(n+3)(n+2)(n+1)(n)(n-1)(n-2)(n-3)} \tag{3.2.6}
\end{aligned}$$

$$\begin{aligned}
& \frac{1}{(n+1)(n+2)(n+3)(n+4)} [[(\alpha_7(n+1)[n^3 + 2n^2 - 135n - 688] + 840\alpha_7) - (\beta_6(n+1)(n^2 + 23n + 102) \\
& - 120\beta_6) - (\gamma_5(n+1)(3n + 16)] - 20\gamma_5) - (2\lambda_4(n - 1) + \mu_3)] a_{n-3} + [(\alpha_6(n+1)(n^3 - 9n^2 - 94n - 282) + \\
& 360\alpha_6) - (\beta_5(n+1)(n^2 + 20n + 276) - 60\beta_5) - \gamma_4(n+1)[3n + 14] - 12\gamma_4) - 2\lambda_3(n - 14) + \mu_2)] a_{n-2} + \\
& [(\alpha_5(n+1)[n^3 + 4n^2 - 59n - 234] - 120\alpha_5) - (\beta_4(n+1)(n^2 + 17n + 54) + 24\beta_4) - \gamma_3(n+1)(3n + 36) + 6\gamma_3) \\
& - 2\lambda_2(n + 2) - \mu_1)] a_{n-1} + [\alpha_4(n+1)[n^3 + 5n^2 - 30n - 120] + 24\alpha_4) - (\beta_3(n+1)[n^2 - 4n - 24] - 6\beta_3) -
\end{aligned}$$



$$(\gamma_2(n+1)(3n+2) - 2\gamma_2) - (\lambda_1(2n+1) + \mu_0)]a_n - \tau_1 C_{n+4}^{n+7} - \tau_2 C_{n+4}^{n+6} - \tau_3 C_{n+4}^{n+5} - \tau_4 C_{n+4}^{n+4}$$

$$= \frac{f_{n-3}}{(n+4)(n+3)(n+2)(n+1)(n)(n-1)(n-2)} \quad (3.2.7)$$

$$\frac{1}{(n+2)(n+3)(n+4)(n+5)} [[(\alpha_7(n+2)[n^3 + 5n^2 - 128n - 822] + 840\alpha_7) - (\beta_6(n+2)(n^2 + 25n + 12)$$

$$- 120\beta_6) - (\gamma_5(n+2)(3n+19)] - 20\gamma_5) - (2\lambda_4 n + \mu_3)]a_{n-2} + [(\alpha_6(n+2)(n^3 + 6n^2 - 85n + 520) +$$

$$360\alpha_6) - (\beta_5(n+2)(n^2 + 22n + 297) - 60\beta_5) - \gamma_4(n+2)(3n-5) - 12\gamma_4) - 2\lambda_3(n-13) + \mu_2)]a_{n-1} +$$

$$[(\alpha_5(n+2)[n^3 + 7n^2 - 48n - 300] - 120\alpha_5) - (\beta_4(n+2)(n^2 + 19n + 72) + 24\beta_4) - \gamma_3(n+2)(3n+15) + 6\gamma_3)$$

$$- 2\lambda_2(n+3) - \mu_1)]a_n - \tau_1 C_{n+5}^{n+7} - \tau_2 C_{n+5}^{n+6} - \tau_3 C_{n+5}^{n+5} = \frac{f_{n-2}}{(n+5)(n+4)(n+3)(n+2)(n+1)(n)(n-1)} \quad (3.2.8)$$

$$\frac{1}{(n+3)(n+4)(n+5)(n+6)} [[(\alpha_7(n+3)[n^3 + 8n^2 - 11n - 944] + 840\alpha_7) - (\beta_6(n+3)(n^2 + 27n + 152)$$

$$- 120\beta_6) - (\gamma_5(n+3)(n+14)] - 20\gamma_5) - (2\lambda_4(n+1) + \mu_3)]a_{n-1} + [(\alpha_6(n+3)(n^3 + 9n^2 + 70n - 600) +$$

$$360\alpha_6) - (\beta_5(n+3)(n^2 + 24n + 320) - 60\beta_5) - \gamma_4(n+3)(3n+20) - 12\gamma_4) - 2\lambda_3(n-12) + \mu_2)]a_n +$$

$$- \tau_1 C_{n+6}^{n+7} - \tau_2 C_{n+6}^{n+6} = \frac{f_{n-1}}{(n+6)(n+5)(n+4)(n+3)(n+2)(n+1)(n)} \quad (3.2.9)$$



$$\begin{aligned}
 & \frac{1}{(n+4)(n+5)(n+6)(n+7)} [[(\alpha_7(n+4)[n^3 + 11n^2 + 58n - 630] + 840\alpha_7) - (\beta_6(n+4) \\
 & (n^2 + 29n + 180)120\beta_6) - (\gamma_5(n+4)(3n+25) - 20\gamma_5) - (2\lambda_4(n+2) + \mu_3)]a_n \\
 & - \tau_1 C_{n+7}^{n+7}] = \frac{f_n}{(n+7)(n+6)(n+5)(n+4)(n+3)(n+2)(n+1)} \quad (3.3.0)
 \end{aligned}$$

A Numerical Experiment

We consider here the following problems for experiment with our results of the preceding sections. The exact error is defined by $\varepsilon * = \max_{0 \leq x \leq 1} [|Y(x_k) - Y_n(x_k)|], 0 \leq x \leq 1, [x_k] = [0.01k], k = 0(1)100$

Example 4.1

$$Ly(x) \equiv y^{iv}(x) + \frac{1}{4}(16 + x^3)y''(x) + x^3y(x) = x^3$$

$$y(0) = 1, y'(0) = 2, y''(0) = 0, y'''(0) = -8$$

Exact solution: $y(x) = 1 + \sin 2x$

$$m = 4, \quad s = 3$$

The linear equations obtained were solved by Maple 18 package.

Table 4.1a: Numerical Results for Example 4.1a(case $n = 7$)


| x | Exact | Approximation ($n = 7$) | Errors |
|----------|--------------|---|---------------|
| .0 | 1.0000000000 | 1.0000000000 | 0.0000e+00 |
| .1 | 1.1986693310 | 1.1986693060 | 2.5190e-08 |
| .2 | 1.3894183420 | 1.3894182070 | 1.3488e-07 |
| .3 | 1.5646424730 | 1.5646424300 | 4.2896e-08 |
| .4 | 1.7173560910 | 1.7173565440 | 4.5325e-07 |
| .5 | 1.8414709850 | 1.8414717560 | 7.7102e-07 |
| .6 | 1.9320390860 | 1.9320390540 | 3.2322e-08 |
| .7 | 1.9854497300 | 1.9854480930 | 1.6373e-06 |
| .8 | 1.9995736030 | 1.9995722210 | 1.3823e-06 |
| .9 | 1.9738476310 | 1.9738500380 | 2.4070e-06 |
| 10 | 1.9092974270 | 1.9092938860 | 3.5406e-06 |

**Table 4.1b: Numerical Result for Example 4.1b($n = 8$)**

| x | Exact | Approximation | Error |
|-----|--------------|---------------|--------------|
| .0 | 1.0000000000 | 1.0000000000 | $0.0000e+00$ |
| .1 | 1.1986693310 | 1.1986693330 | $2.0832e-09$ |
| .2 | 1.3894183420 | 1.3894183460 | $3.4859e-09$ |
| .3 | 1.5646424730 | 1.5646424580 | $1.5186e-08$ |
| .4 | 1.7178560910 | 1.7173560680 | $2.2947e-08$ |
| .5 | 1.8414709850 | 1.8414710090 | $2.3801e-08$ |
| .6 | 1.9320390860 | 1.9320391540 | $6.8033e-08$ |
| .7 | 1.9854497300 | 1.9854497230 | $6.9754e-09$ |
| .8 | 1.9995736030 | 1.9995734820 | $1.2100e-07$ |
| .9 | 1.9738476310 | 1.9738476860 | $5.5465e-08$ |
| 10 | 1.9092974270 | 1.9092972460 | $1.8117e-07$ |

Table 4.1c: Numerical Result for Example 4.1c ($n = 9$)

| x | Exact | Approximation ($n = 9$) | Error |
|-----|--------------|---------------------------|--------------|
| .0 | 1.0000000000 | 1.0000000000 | $0.0000e+00$ |
| .1 | 1.1986693310 | 1.1986693310 | $9.1858e-11$ |
| .2 | 1.3894183420 | 1.3894183420 | $1.1671e-10$ |
| .3 | 1.5646424730 | 1.5646424730 | $7.2551e-10$ |
| .4 | 1.7173560910 | 1.7173560910 | $5.7954e-10$ |
| .5 | 1.8414709850 | 1.8414709870 | $2.1114e-09$ |
| .6 | 1.9320390860 | 1.9320390850 | $1.0526e-09$ |
| .7 | 1.9854497300 | 1.9854497260 | $4.1769e-09$ |
| .8 | 1.9995736030 | 1.9995736060 | $3.3152e-09$ |
| .9 | 1.9738476310 | 1.9738476330 | $2.4362e-09$ |
| 10 | 1.9092974270 | 1.9092974360 | $9.2127e-09$ |



CONCLUSION

The derivation of an approximate scheme for a fourth order differentiation equations with third degree overdetermination by the integrated formulation of the tau method has been presented. The class of ordinary differential equations under consideration were integrated four times and then pertubing the resulting equation. This is to garantte an improved accuracy of the desired approximation viz-a-viz those of the recursive and the differential formulations. Numerical evidences from the application of the Integration Scheme show that it is accurate and effective.

REFERENCES

- [1] Lanczos, C. (1983): Trigonometry Interpolation of Empirical and Analytical Function. Math and Physics, 17, 123-199.
- [2] Onumanyi, P. and Ortiz, E.L (1982):Numerical Solution of Higher Order Boundary Value Problems for Ordinary Differnetial Equations with an Estimation of the Error, Intern. J. Numer. Mech. Engrg, 18, 775-781.



- [3] Ortiz E.L. (1974): Canonical Polynomial in the Lanczos Tau method, Studies in Numerical Analysis (edited by Scaife B.K.P), Academic Press, New York.
- [4] Fox, L. and Parker, I.B (1968): Chebyshev Polynomials in Numerical Analysis, Oxford University Press Oxford
- [5] Mousavi S. and Mousavi W. (2012) New Application of Tau Method for Solving Differential Equations of the Ian Emden type. World Applied Science Journal 18(5): 642-644.
- [6] Adeniyi R.B. (1991). On The Method of Tau Method for Numerical Solution of Ordinary Differential Equations. Doctoral Thesis, University of Ilorin, Ilorin, Nigeria.
- [7] Aliyu I.M.A (2012). Computational Error Estimate of Integrated Formulation of the Tau Method for some class of ODEs. Doctoral Thesis, University of Ilorin, Ilorin, Nigeria.
- [8] Lanczos, C. (1956): Applied Analysis, Prentice Hall, New Jersey.
- [9] Adeniyi R.B. and Aliyu A.I. (2011). Malli on the Integrated Formulation of the Tau Method involving at most Two Parameters. International Journal on Research and Development, Book of proceedings 3:104-114.
- [10] Yisa B.M. (2012). The recursive formulation of the Tau method for non-over determined ODEs, Doctoral Thesis, University of Ilorin, Ilorin, Nigeria.
- [11] Adeniyi R.B. and Aliyu A.I. (2007). Maali on the integrated formulation of numerical schemes for a class of Non-overdetermined second order Differential Equations. Nigeria Mathematical Society, 29:111-128.
- [12] Ojo ,V.O. and Adeniyi R.B.(2012).The differential form of the Tau Method and its Error estimate for Third Order Non-Overdetermined Differential Equations.General Mathematics Notes, 11(1):41-49.
- [13] Ojo, V.O. and Adeniyi R.B.(2012).The Integrated Formulation of the Tau Method and its Error Estimate for Third Order Non-Overdetermined Differential Equations. International Journal of Engineering Research and Technology (IJERT).ISSN 2278-0181, 1 (9).