



DETERMINATION OF THE DEGREE OF HOMOGENEITY OF PRIMITIVE PERMUTATION GROUPS VIA THE SOCLE OF THE GROUPS

Danbaba Adamu* and Momoh Umoru Sunday

Department of Mathematics University of Jos, Jos Nigeria.

*Corresponding Author's Email: adamu0019@gmail.com

Cite this article:

Danbaba, A., Momoh, S. U. (2024), Determination of the Degree of Homogeneity of Primitive Permutation Groups via the Socle of the Groups. African Journal of Mathematics and Statistics Studies 7(3), 190-197. DOI: 10.52589/AJMSS-84OWWYYB

Manuscript History

Received: 21 Jun 2024

Accepted: 28 Aug 2024

Published: 9 Sep 2024

Copyright © 2024 The Author(s).

This is an Open Access article distributed under the terms of Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0), which permits anyone to share, use, reproduce and redistribute in any medium, provided the original author and source are credited.

ABSTRACT: *The nilpotency class for the Frobenius was determined based on the structure theorem. The socle of the groups were observed to be regular normal and elementary abelian such features were the conditions for the nilpotency classes, as they were the basis on which the socle of these groups constructed were nilpotent of some classes or order. The socle of the nilpotent groups whose structures is in conformity with D were classified based on the classification scheme for the finite primitive groups in relation to socle type.*

KEYWORDS: Frobenius, groups. Socle, nilpotency, finite, abelian, regular.



INTRODUCTION

The structure of a finite Frobenius group was as a result of the work carried out on primitive groups by [11] and was further investigated by Burnside [2]. He showed that a finite Frobenius group has proper nontrivial characteristic subgroups. Therefore, this idea played a key role in the study of finite 2-transitive permutation groups. The proof of the work of [11] was geared toward the use of character theory. The major work on this group was carried out by [2]. He gave the outline of the structure for finite Frobenius but in the real sense, the structure of the Frobenius group was described by Frobenius in paper [11], Zassenhaus and Thompson as stated in [10].

Let G be a finite Frobenius group and with point stabilizer such that;

$$(D) \quad K = \{ x \in G \mid x = 1 \text{ or } \text{fix}(x) = \emptyset \}$$

Then D is a subgroup of G and is nilpotent, also for any prime, p , for which the Sylow p -subgroup of G has a point stabilizer which is cyclic. Throughout we denote D as the Frobenius subgroup.

The statement for the structure of finite Frobenius groups as given in D was due to work of the work of Frobenius in [11] which was later generalised in the work of [18]. It follows that a permutation group which is a Frobenius group based on the above conditions has a regular abelian subgroup and the point stabilizer has only one element of order 2. He further state that if H is a primitive group but not regular and, also $\frac{3}{2}$ -transitive, then H is normal in G ,

Therefore, every finite primitive group has a unique minimal normal subgroup which is regular abelian and simple and also isomorphic to each other. Also [13] worked on finite p - groups with a Frobenius group of automorphism, whose kernel is a cyclic p - group. In his work, he defined a Frobenius group as a finite normal subgroup which is non-trivial. We intend in this paper to determine the socle of primitive with a subgroup satisfying the condition defined in D, with a view to determine subgroup generated by minimal normal subgroups with a structure satisfying D that is the socle of the group. The classification scheme for finite primitive based on the O'Nan-Scott theorem rest basically on the socle type of the finite primitive groups.

Preliminary result

Theorem 2.1: Let G be a finite group which is not regular and H a subgroup with G_α a point stabilizer of G . Therefore G can be written as $G = HG_\alpha$.

Theorem 2.2: Let G be a finite Frobenius group and G_α be a point stabilizer and let D be defined, then the following hold

1. D is a subgroup of G which is normal and regular in G
2. For each odd prime, p , the Sylow p -subgroup of G_α is cyclic.
3. D is nilpotent

The structure theorem for the Frobenius group is the necessary and sufficient condition for a subgroup to have a structure of D of a Frobenius group. The subgroup are fix point free.



Suppose G is transitive then D is a derangement except for the identity subgroup. With these we take the following.

Remark 2.3:

In the case that G_α is not soluble, then G has only one abelian composition factor which is A_5

We now state a theorem which gives us a condition for the existence of a Frobenius group with a point stabilizer having only one element of even order. This assertion only satisfy the condition in which $p = 2$ exactly, since it is the only prime number which is also even.

The next result is given in the relation to the concept of reugular abelian groups defined in [14]

Theorem 2.4: Let G be a finite Frobenius group of degree n , and D a subgroup. If G_α has even order then D is a regular normal abelian subgroup of G and G_α has exactly one element of order 2.

Proof:

For G_α to have even order it imply that it have elements of order of two. Let T be the G –conjugacy class containing these elements. Since the point stabilizers are disjoint, each of the point stabilizer contains at least one element from T and $|T| \geq n$. Consider the cycle decomposition of an element $t \in T$ such that t has one cycle of length 1 and $\frac{n-2}{2}$ –cycle of length 2. since no nontrivial element of G has more than one fixe point, no two elements from G can contain the same 2 –cycle. There are exactly $\frac{n(n-1)}{2}$, 2 –cycles in $\text{sym}(\Omega)$, and so we conclude that $\frac{|T|(n-1)}{2} \leq \frac{n(n-1)}{2}$ and hence $|T| \leq n$. But $|T| \geq n$, therefore $|T| = n$, and every 2 –cycle occurs in one of the element of T . In particular, each point stabilizer contains exactly one element from T , and T contains all elements of order 2 in G .

We suppose that $st \in D$ for any $s, t \in T$ then we assume that $\text{fix}(st) = \Phi$ for if not then any $\beta \in \Omega$ we have $\text{fix}(st) = \beta$ which contry to the definition in D and we may have $\beta^t = \beta^{(st)t} = \beta$ for any distinct s and t , and so eithr $(\beta \beta^s) = (\beta \beta^t)$ is a 2 –cycle in both s and t . But the case is not possible. Therefore $st \in D$ as asserted.

$\text{fix}(t) \in T$ then $tT \subseteq D$, and since both have size n we onclude that $tT = D$ in particular $1 \in D$ and therefore, $DD^{-1} \subseteq TT \subseteq D$, and so D is a subgroup and further D is abelian .

The next result is due to the idea of a statement of result in [10].

Theorem 2.5: Let G be a 2 –transitive Frobenius group, and D a subgroup. Then suppose that either G is finite or the point stabilizer G_α is abelian, then D is regular normal abelian subgroup of G in which each nontrivial element has same order.

Proof:

Suppose $|\Omega| = n$ then $|D| = n$ for if G is 2 –transitive then $|G_\alpha|$ divides $n - 1$ therefore G is a Frobenius group if for any $u \neq 1$ and $u \in D$ then $C_G(u) \subseteq D$ then $|G : C_G(u)| \geq n - 1$. Therefore u has atleast one conjugate element in G . On the other hand each conjugate element of u is clearly a nontrivial element from D then we conclude that $C_G(u) = D$. thus we have



shown that D is a subgroup and each element of D lies in the centre of D . Hence D is elementary abelian p -group and so D is regular and normal. Further based on Theorem 2.4 D is abelian also since G is Frobenius then for any $s, t \in D$ and $\alpha, \beta \in \Omega$ then $(\alpha\beta)^t = (\beta\alpha)$, therefore, for any $(t \neq 1)$, it implies $t^2 = 1$, and so t fixes both α and β . Conversely, if G is Frobenius then every element $x, y \in G$ implies $x, y^{-1} \in D$ and so G is of prime degree, and so by Theorem 2.4, D is nilpotent. The statement also follows from Theorem 2.2, as such D is cyclic which by implication D is abelian. Further for any $u, v \in D$ there is $\alpha, \beta \in \Omega$ such that $\alpha^u = \beta^v = 1$ and so $u = v$. Hence the theorem.

Theorem 2.6 [10]: Let G be a finite primitive group with abelian point stabilizer, then G is either,

regular or of prime degree or a Frobenius group.

Proof: Since G is a primitive group and not regular, then G is a Frobenius group and has a subgroup which is a Sylow p -subgroup, which is regular and abelian. Therefore G_α is of the type D , and so D is nilpotent which also by Theorem 2.2, it shows condition 1 implies condition 3 thus G is Frobenius

Conversely, suppose G is Frobenius, it follows from Theorem 2.1 that, G has a subgroup which is regular and abelian of prime degree, therefore the subgroup may be of type D . Hence G has G_α as the only proper subgroup and so G is primitive.

Next we define nilpotency of a finite primitive group as it is a requirement in the attainment of a Frobenius structure.

Definition 2.7: A finite primitive group G is nilpotent if and only if G has an upper central and a lower central series. That is there is an integer $k \geq 0$ such that $\gamma_k(G) = 1$.

Therefore we state categorically that if G is nilpotent and N is normal then G/N is nilpotent. Also if G is nilpotent then it implies G is soluble. The next result clearly gives condition for the existence of nilpotency classes

Theorem 2.8: Let G be a finite primitive group with a nilpotent point stabilizer. Then G is soluble if the Sylow 2-subgroup of the point stabilizer is also nilpotent of class at most 2.

Proof: Suppose G has a nilpotent point stabilizer, then G_α is cyclic and so the point stabilizer is an elementary p -abelian group. Therefore by Definition 2.6 and the condition that G_α is a normal subgroup of G it implies that G is soluble which follows immediately from Theorem 2.2.

Conversely if G is of order say, np^a for $a \geq 1$ then G contains a normal subgroup which is a Sylow p -subgroup. Therefore it can be deduced from Theorem 2.1 that G_α is nilpotent, and so by Theorem 2.2(2) G_α is nilpotent of class at most 2.

Theorem 2.9: Let G be a finite primitive group with a maximal subgroup M which is abelian, then G is solvable.

Proof: If G is primitive then for any normal subgroup of G , say D is maximal implying $M \leq D$, so we assume $D = M$. Therefore the composition series for G is of the form $G \triangleright D \triangleright 1$ and



so D is a normal subgroup of G . Moreover since G is primitive it imply that D is abelian and so it is of order a power of p .

Remark 2.10: the necessary and sufficient condition that a finite primitive group is nilpotent is G is a finite p –group and abelian.

Theorem 2.11: Let G is p –group then G is nilpotent

Proof: Suppose $|G| = 1$, the result is trivially true. We assume that $|G| \geq 2$. Also assume inductively that the theorem holds for all p –groups of order less than $|G|$ hence it iply $G/Z(G)$ is nilpotent. Thus G is nilpotent if and only if $G/Z(G)$ is cyclic.

Therefore we take the statement of the following theorem whose proof will not be given here, it is based on the work carried out in [1]

Theorem 2.12 [1]: Let G be a primitive group. The following are equivalent

- i. G is nilpotent
- ii. Every subgroup of G is subnormal in G
- iii. Whenever H a proper subgroup of is G then H is a proper subgroup of its normalizer in G .
- iv. Every maximal subgroup of G is normal in G
- v. $G' \leq \vartheta(G)$
- vi. Every Sylow p –subgroup of G is normal in G
- vii. G is a direct product of groups of prime power order.

The next theorem is as a result of the work of [17] in relation to the structure of D which ill help in determining the socle for G having a subgroup in the form of D .

Theorem 2.13 [10]: Let G be a group which acts primitively and on Ω with $|\Omega| = n$. Let $H = \text{soc}(G)$ and $\alpha \in \Omega$. Then H is of type T then G is affine and T is abelian of order p and $n = p^n$ and G_α is a complement to which acts on H and is simple.

We can say therefore, if G has a minimal normal subgroup K say, then for some prime p and some integer (d, k) , G is a regular elementary abelian group of order p^d and $\text{soc}(G) = K = C_G(K)$.we observe further, if K has the structure of D the $\text{soc}(G) = D = C_G(D)$ which is also an elementary abelian group of prime power p^d and also isomorphic to an affine group. This suffices to say D is a subgroup and is nilpotent. Thus we can clearly state that socle are subgroups of normalizers subgroups

Theorem 2.1 [10] 4: Let G be a finite group, then $\text{soc}(G) \leq N_G(H)$ for each subnormal group H of G .



Proof:

The result is certainly true if $G = H$. Therefore induction on $|G:H|$ with $H < G$ show that each minimal normal subgroup K of G is contained in the normalizer of $N_G(H)$.

Since it is subnormal, there exist $L \triangleleft G$ with $H \leq L \triangleleft G$. Then either $K \leq L$, $(K, L) = KL$. Hence $K \leq C_G(H) \leq N_G(H)$ and the result is true. So suppose that $K \not\leq L$. Then there exists a minimal normal subgroup of $x^{-1}L = L$, and induction shows that

$$K = \{x^{-1}Hx\} \leq N_L(H) \leq N_G(H).$$

Since K is a minimal normal subgroup of G we have that

$$K = \{x^{-1}Hx \mid x \in G\}. \text{ Implying } K \leq N_G(H).$$

MAIN RESULTS

Theorem 3.1 : Let G be a finite group of order n and H a subgroup of order p^a for $a \geq 1$ then $\text{soc}(G)$ is a regular abelian group of order a power of p .

Proof:

For G to regular imply G then by Theorem 2.6 show that G is either a Frobenius group or regular if and only if G has an abelian stabilizer so let H be a subgroup of G with the structure as defined in D . If $H = G_\alpha$ then H is regular and abelian, also suppose H and maximal with $H = \text{soc}(G)$ imply H is nilpotent.

Conversely suppose G is nilpotent and H is normal in G then every Sylow 2 -subgroup of G is an elementary p -abelian group of order a power of p . Therefore if H is maximal Theorem 2.13 imply $\text{soc}(G)$ is regular and abelian of order p .

Theorem 3.2: Let G be a finite nilpotent group with a regular normal subgroup H of prime order. Then $\text{soc}(G)$ is nilpotent.

Proof: Since G is nilpotent it shows that the normal subgroups of G is abelian and there let H be a normal of G with the structure as defined in D then H is regular and abelian and is also maximal, otherwise there may be a chain of subgroup with another proper subgroup say, M such that $G \supset H \supset M \supset \{e\}$ for $M \leq H$, this shows that G is nilpotent of class 3. But H is a maximal subgroup of G therefore it is nilpotent of class at most 2 by Theorem 2.7, and so, suppose $H = G_\alpha$ and that $\text{soc}(G) = H$ then by theorem 2.12 G is an elementary p -abelian group which imply $\text{soc}(G)$ is nilpotent.

Theorem 3.3: Let G be a finite nilpotent group and H a subnormal group of G , therefore $\text{soc}(G)$ is an elementary abelian p -group

Proof: suppose G is primitive of degree n , imply that $|G| = n = p^a m$, and G has Sylow subgroup by first sylows theorem. Let H be the subgroup of G , since G is primitive show that H is maximal and so by Theorem 3.3, H is of type D above consequently $H = \text{soc}(G)$ which is also elementary p -abelian



Conversely if G is an abelian p -group, it implies that any chain of subgroups of G has a maximal subgroup H . Say, therefore by Theorem 2.7 it implies that $H = \text{soc}(G)$ and so H is nilpotent of class 2 and so by Theorem 3.3, the chain $G \supset H \supset M \supset \{e\}$ has a subnormal group M in line with Theorem 2.14 and so $M = H$, hence G is nilpotent.

CONCLUSION

The regular subgroups of the type as defined in D and their nilpotency classes were obtained for Frobenius groups with regular normal abelian subgroups. These groups were of order a power of p in which most of the subgroups had the structure of D . The socle of G had a direct relation with the classification scheme for finite simple groups based on the socle type as is in the work of [17]. The case for which p was of order 2 was determined for groups of even order. It showed that the socle of the groups having the structure of D were transitive and nilpotent.

Abbreviations

$\text{soc}(G)$ - Socle of the group G

$\text{fix}(x)$ - Fix of an element x in G

$\text{Supp}(x)$ - Support of the element x in G

CONFLICTS OF INTEREST

There are no conflicts of interest except for lack of funds, which will serve as a catalyst for innovation and the development of the field of group theory.

REFERENCES

- [1] Audu, M. S., & Momoh, S.U (1990): On transitive Permutation groups. *Abacus*, 19,(2). 17-23.
- [2] Burnside, W (1911). *Theory of groups of finite order*. Cambridge University press. London.
- [3] Cameron,P.J. (1981). Finite permutation groups and finite simple groups; *London Math. Soc.* DOI 101112/BLMS/13.1.1.
- [4] Cameron, P.J. (1999). Permutation groups. *Math. Soc. Student Text 45*, Cambridge University Press.
- [5] Cameron,P.J. (2000). Aspect of infinite permutation groups: School of Mathematical Sciences, Queen Mary, *University of London*.
- [6] Cameron, P.J. (2000). t -orbit homogenous permutations. *London Math Soc. Subject Classification 20B10*, DOI:10.1112/Sooooo0000000000000000.
- [7] Cameron,P.J. (2004). The encyclopaedia of design theory on primitive permutation groups. www.maths.qmul.ac.uk. Retrieved on 12/07/2015.
- [8] Cameron,P.J (2012). Permutation groups and regular semi groups. *London Math Subject classification*.1273-1285



-
- [9] Cameron, P.J. (2013) Permutation group and transformation semi-group, Novi sad Algebraic Conference. *University of St. Andrews*
- [10] Dixon, J., & Mortimer, B (1996). *Permutation groups Graduate texts in mathematics*. Springer New York`.
- [11] Frobenius ,F.G. (1904). Uber die charaktere der mehch trasitiven gruppen.mouton de grupter 558-571. *Berlin*.
- [12] Kantor, W. M. (1972). k -Homogeneous permutation groups. *Math Z*, 124, 261-265 .*Springer Verlag*.
- [13] Khukhr, O (2013). Finite p -groups with a Frobenius group of automorphism whose kernel is cyclic p -group. *Manchester institute for Mathematical Sciences: School of Mathematics, University of Manchester*.
- [14] Liebeck, W., & Praeger C.E, & Sax, I .J. (2000). Primitive permutation groups with a regular subgroup. *Journal of Mathematics subject classification*. 20B15, 05C25.
- [15] Livingstone, D., & Wagner, A. (1965). Transitivity of finite permutation groups on unordered sets. *Math Z*. 90 393-403.
- [16] Martin, W.J., & Segan, B (2000). New notion of transitivity for groups and set of permutations. *Journal of Math Soc. Math Subject Classification 20B20*.
- [17] O'Nan ,M.E., & Scott, L (1979). Finite groups. Santa Cruz conference. *London Math Soc* (2)32.
- [18] Wielandt, H. (1969). Permutation groups through the invariant relation and invariant functions. *Columbus*.399514.
- .
- .