



ON THE EXISTENCE OF MAXIMAL SOLUTION AND LYAPUNOV PRACTICAL STABILITY OF NONLINEAR IMPULSIVE CAPUTO FRACTIONAL DERIVATIVE VIA COMPARISON PRINCIPLE

Jackson Efiang Ante^{1*}, Samuel Okon Essang², Okoi Okoi Itam³, and Etorti Imoke John⁴

¹Department of Mathematics, Topfaith University, Mkpatak, Nigeria.

Email: jackson.ante@topfaith.edu.ng

²Department of Mathematics, Arthur Jarvis University, Akpabuyo, Calabar, Nigeria.

Email: sammykmf@gmail.com

³Department of Mathematics, University of Calabar, Calabar.

Email: itamokoi@gmail.com

⁴Department of Mathematics, Cross River State College of Education, Akamkpa, Nigeria.

Email: imoke.etorti@crs-coeakamkpa.edu.ng

*Corresponding Author's Email: jackson.ante@topfaith.edu.ng

Cite this article:

Ante, J. E., Essang, S. O., Itam, O. O., John, E. I. (2024), On the Existence of Maximal Solution and Lyapunov Practical Stability of Nonlinear Impulsive Caputo Fractional Derivative via Comparison Principle. Advanced Journal of Science, Technology and Engineering 4(4), 92-110. DOI: 10.52589/AJSTE-9BWUJX90

Manuscript History

Received: 14 Aug 2024

Accepted: 17 Oct 2024

Published: 29 Oct 2024

Copyright © 2024 The Author(s). This is an Open Access article distributed under the terms of Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0), which permits anyone to share, use, reproduce and redistribute in any medium, provided the original author and source are credited.

ABSTRACT: *This paper examines the existence of maximal solution of the comparison differential system and also establishes sufficient conditions for the practical stability of the trivial solution of a nonlinear impulsive Caputo fractional differential equations with fixed moments of impulse using the vector Lyapunov functions. First, it was discovered that the vector form of the Lyapunov function was majorized by the maximal solution of the comparison system. From the results obtained, it was established that the main system is practically stable in the sense of Lyapunov.*

KEYWORDS: practical stability, Caputo derivative, impulse, vector Lyapunov functions, fractional differential equations, maximal solution.



INTRODUCTION

The history of fractional calculus has been about 300 years old, and its development is mainly focused on the pure mathematical field [27]. The earliest – more or less systematic studies of the concept, seem to have been made in the 19th century by Liouville, Riemann, Leibnitz, Caputo, etc. [37], and in the last four decades, it was observed that the advancement of fractional calculus has enabled the description of complex system behaviors through fractional differential systems (including many physical phenomenon having memory and genetic characteristics), thus providing new insights into their dynamics.

One of the trends in the stability theory of solutions of differential equations is the so-called practical stability [8, 27, 31, 32, 33]. This aspect of stability was introduced by [24] and it is used in estimating the worst-case transient and steady-state responses together with verifying point-wise in time constraints imposed on the solution path or the trajectory curve.. Fundamental results have been obtained for fractional order derivative using the auxiliary Lyapunov's functions which are analogues of vector Lyapunov functions, by means of the comparison method.

Alongside the development of the theory of practical stability in recent years is the mathematical theory of impulsive differential equations which have experienced a massive research attention and development. Now, the theory of impulsive differential equations is richer than the corresponding theory of differential equations [16] as they constitute very important models for describing the true state of several real life processes and phenomena since many evolution processes are characterized by the fact that, at certain moments of time, they experience a change of state abruptly. These processes are assumed to be subject to short term perturbations whose duration is negligible in comparison with the duration of the process. Consequently, it is natural to assume that these perturbations act instantaneously, that is, in the form of impulses. For instance, many biological phenomena involving thresholds, bursting rhythm models in medicine and biology, optimal control models in economics, pharmacokinetics and frequency modulated systems do exhibit impulsive effects [16].

Moreover, the efficient applications of impulsive differential system require the finding of criteria for stability of their solutions [35], and one of the most versatile methods in the study of the stability properties of impulsive systems is the Lyapunov function (Lyapunov second method) The method was originally developed for studying the stability of a fixed point of an autonomous or nonautonomous differential equations. However, as was argued in [29], it was then extended from fixed points to sets, from differential equation to dynamical systems and to stochastic equations.

Suffice to say that the novelty of the Lyapunov's second method over other methods of examining stability properties of impulsive differential systems like the Razumikhin technique, the use of matrix inequality, etc. stems from the fact that the method allows us to examine the stability of solutions without first solving the given differential equation by seeking an appropriate continuously differentiable function (called Lyapunov function) that is positive definite and whose time derivative along the trajectory curve is negative semidefinite.



The stability of the zero solution of impulsive differential equations have been extensively studied in [11,30]. Furthermore, the study of stability of fractional order systems is quite recent and one of the main difficulties in the application of the Lyapunov function to fractional order differential equations is the appropriate definition of its derivative among the fractional differential equations. Thus, to allay this problem, [1] adopted the choice of a Lyapunov function that is continuously differential, with a re-definition of the Dini derivative for the given fractional order system. This choice was necessary because, the choice of the Dini derivative for the fractional differential equation as was used in [17-18] had some “restrictions and difficulties” (see [1] and the references therein).

Now, the stability of fractional order systems using the scalar Lyapunov function have been examined in [1-2,5,9,41]. Using the generalized Caputo fractional Dini derivative and scalar impulsive fractional differential equations, [2] established the comparison results together with sufficient conditions for the stability properties of impulsive fractional differential equations. However, the set-back in this approach arise from the fact that, when the system becomes complex or large, the scalar Lyapunov functions lacks a definite algorithm for handling such systems. Due to this pitfall, the use of vector Lyapunov functions becomes very necessary because of its ability to handle complex systems as well as large scale dynamical systems. The method of vector Lyapunov function involves splitting the Lyapunov functions into several components, so that each of the components can adequately describe the system state. In this way, Lyapunov functions are easily constructed, and the conditions ensuring the required stability are less restrictive (see [40] and the references therein). Fundamental results for the stability of impulsive Caputo fractional differential equations have been examined in [1] and [2].

In this paper, the existence of maximal solution of the comparison system for vector Lyapunov function is established. Again, sufficient condition for the practical stability of impulsive Caputo fractional order systems is presented by means of the comparison principle. It was discovered that the vector form of the Lyapunov function is majorized by the maximal solution of the comparison system. The sufficient conditions for the practical stability of impulsive fractional order systems as established stressed the fact that the solution of the main system is practically stable.

Preliminary Notes and Definitions

Fractional calculus is seen as a natural generalization of the classical calculus of integer order and thereby allows for the extension of the traditional concepts of derivative and integral to functions with fractional orders. By this extension, functions with noninteger orders are much more flexible in describing real world systems (see [14,25,26,34,37]).

There are several definitions of fractional derivatives and fractional integrals.

General case. Let the number $n-1 < \beta < n, \beta > 0$ be given, where n is a natural number, and $\Gamma(\cdot)$ denotes the Gamma function.

Definition 2.1.

The Riemann Liouville fractional derivative of order β of $\gamma(t)$ is given by (see [35])



$${}^{RL}D_t^\beta \gamma(t) = \frac{1}{\Gamma(n-\beta)} \frac{d^n}{dt^n} \int_{t_0}^t (t-s)^{n-\beta-1} \gamma(s) ds, t \geq t_0$$

Definition 2.2.

The Caputo fractional derivative of order β of $\gamma(t)$ is given by (see [35])

$${}^C D_t^\beta \gamma(t) = \frac{1}{\Gamma(n-\beta)} \int_{t_0}^t (t-s)^{n-\beta-1} \gamma^{(n)}(s) ds, t \geq t_0$$

The Caputo derivative has many properties that are similar to those of the standard derivatives which make them easier to understand and apply. Also, the initial conditions of the Caputo fractional order derivative are also easier to interpret in physical context.

Definition 2.3.

The Grunwald-Letnikov fractional derivative of order β of $\gamma(t)$ is given by (See [1])

$${}^{GL}D_0^\beta \gamma(t) = \lim_{h \rightarrow 0^+} \frac{1}{h^\beta} \sum_{r=0}^{\lfloor \frac{t-t_0}{h} \rfloor} (-1)^r {}^\beta C_r \gamma(t-rh), t \geq t_0$$

and

Definition 2.4. The Grunwald-Letnikov fractional Dini derivative of order β of $\gamma(t)$ is given by (See [1])

$${}^{GL}D_0^\beta \gamma(t) = \limsup_{h \rightarrow 0^+} \frac{1}{h^\beta} \sum_{r=0}^{\lfloor \frac{t-t_0}{h} \rfloor} (-1)^r {}^\beta C_r \gamma(t-rh), t \geq t_0$$

where ${}^\beta C_r$ are the binomial coefficients and $\lfloor \frac{t-t_0}{h} \rfloor$ denotes the integer part of $\frac{t-t_0}{h}$.

Particular case (when $n=1$). In most applications, the order of β is often less than 1, so that $\beta \in (0,1)$. For simplicity of notation, we will use ${}^C D^\beta$ instead of ${}^C D_0^\beta$ and the Caputo fractional derivative of order β of the function $\gamma(t)$ is

$${}^C D^\beta \gamma(t) = \frac{1}{\Gamma(\beta)} \int_{t_0}^t (t-s)^{-\beta} \gamma' ds, t \geq t_0 \quad (2.1)$$

3. Impulses in fractional differential equations

Consider the initial value problem (IVP) for the system of fractional differential equations (FrDE) with a Caputo derivative for $0 < \beta < 1$.



$$\begin{aligned} {}^C D^\beta \gamma(t) &= f(t, \gamma), \quad t \geq t_0, \\ \gamma(t_0) &= \gamma_0, \end{aligned} \quad (3.1)$$

where $\gamma \in R^N$, $f \in C[R_+ \times R^N, R^N]$, $f(t, 0) \equiv 0$ and $(t_0, x_0) \in R_+ \times R^N$.

Some sufficient conditions for the existence of the global solutions to (3.1) are considered in [7,11,22, 23,29,35,42].

The IVP for FrDE (3.1) is equivalent to the following Volterra integral equation (See [2]),

$$\gamma(t) = \gamma_0 + \frac{1}{\Gamma(\beta)} \int_{t_0}^t (t-s)^{\beta-1} f(s, \gamma(s)) ds, \quad t \geq t_0 \quad (3.2)$$

Consider the IVP for the system of impulsive fractional differential equations (IFrDE) with a Caputo derivative for $0 < \beta < 1$,

$$\begin{aligned} {}^C D^\beta \gamma(t) &= f(t, \gamma), \quad t \geq t_0, \quad t \neq t_k, \quad k = 1, 2, \dots \\ \Delta \gamma &= I_k(\gamma(t_k)), \quad t = t_k, \quad k \in N, \\ \gamma(t_0^+) &= \gamma_0, \end{aligned} \quad (3.3)$$

where $\gamma, \gamma_0 \in R^N$, $f \in C[R_+ \times R^N, R^N]$, and $t_0 \in R_+$, $I_k : R^N \rightarrow R^N$, $k = 1, 2, \dots$

under the following assumptions:

(i) $0 < t_1 < t_2 < \dots < t_k < \dots$, and $t_k \rightarrow \infty$ as $k \rightarrow \infty$;

(ii) $f : R_+ \times R^N \rightarrow R^N$ is piecewise continuous in $(t_{k-1}, t_k]$ and for each $x \in R^N$, $k = 1, 2, \dots$, and $\lim_{(t,y) \rightarrow (t_k^+, \gamma)} f(t, y) = f(t_k^+, \gamma)$ exists;

(iii) $I_k \times R^N \rightarrow R^N$

In this paper, we assume that $f(t, 0) \equiv 0$, $I_k(0) \equiv 0$ for all k so that we have trivial solution for (3.3), and the points t_k , $k = 1, 2, \dots$ are fixed such that $t_1 < t_2 < \dots$ and $\lim_{k \rightarrow \infty} t_k = \infty$. The system (3.3) with initial condition $\gamma(t_0) = \gamma_0$ is assumed to have a solution $\gamma(t; t_0, \gamma_0) \in PC^\beta([t_0, \infty), R^N)$.

Remark 3.1. The second equation in (3.3) is called the impulsive condition, and the function $I_k(\gamma(t_k))$ gives the amount of jump of the solution at the point t_k .

Definition 3.1. Let $\Omega : R_+ \times R^N \rightarrow R^N$ Then Ω is said to belong to class χ if,

(i) Ω is continuous in $(t_{k-1}, t_k]$ and for each $\gamma \in R^N$ and $\lim_{(t,y) \rightarrow (t_k^+, \gamma)} \Omega(t, y) = \Omega(t_k^+, \gamma)$ exists;



(ii) Ω is locally Lipschitz with respect to its second argument x and $\Omega(t,0) \equiv 0$

Now, for any function $\Omega(t, \gamma) \in PC([t_0, \infty) \times \xi, R_+^N)$. we define the Caputo fractional Dini derivative as:

$${}^c D_+^\beta \Omega(t, \gamma) = \limsup_{h \rightarrow 0^+} \frac{1}{h^\beta} \left\{ \Omega(t, \gamma) - \Omega(t_0, \gamma_0) - \sum_{r=1}^{\lfloor \frac{t-t_0}{h} \rfloor} (-1)^{r+1} C_r [\Omega(t-rh, \gamma - h^\beta f(t, \gamma)) - \Omega(t_0, \gamma_0)] \right\} \tag{3.4}$$

$t \geq t_0$ where $t \in [t_0, \infty)$, $\gamma, \gamma_0 \in \xi$, $\xi \in R^N$ and there exists $h > 0$ such that $t - rh \in [t_0, T]$.

Definition 3.2. A function $g \in PC[R^n, R^n]$ is said to be quasimonotone nondecreasing in γ , if $\gamma \leq y$ and $\gamma_i = y_i$ for $1 \leq i \leq n$ implies $g_i(\gamma) = g_i(y)$.

Definition 3.3. The zero solution of (3.3) is said to be:

(PS1) practically stable if for every $\varepsilon > 0$ and $t_0 \in R_+$ there exist $\delta = \delta(\varepsilon, t_0) > 0$ continuous in t_0 such that for any $\gamma_0 \in R^N$, $\|\gamma_0\| \leq \delta$ implies $\gamma_0 \in R^N$ $\|\gamma(t, t_0, \gamma_0)\| < \varepsilon$ for $t \geq t_0$;

(PS2) uniformly practically stable if for every $\varepsilon > 0$ and $t_0 \in R_+$ there exist $\delta = \delta(\varepsilon) > 0$, continuous in t_0 such that for any $\gamma_0 \in R^N$, $\|\gamma_0\| \leq \delta$ implies $\gamma_0 \in R^N$ $\|\gamma(t, t_0, \gamma_0)\| < \varepsilon$ for $t \geq t_0$;

(PS3) asymptotically practically stable if it is stable and if for each $\varepsilon > 0$ and $t_0 \in R_+$ there exist positive numbers $\delta_0 = \delta_0(t_0) > 0$ and $T = T(t_0, \varepsilon)$ such that for $t \geq t_0 + T$ and $\|\gamma_0\| \leq \delta$ implies $\|\gamma(t, t_0, \gamma_0)\| < \varepsilon$;

(PS4) uniformly asymptotically practically stable if it is uniformly stable and $\delta_0 = \delta_0(\varepsilon)$ and $T = T(\varepsilon)$ such that for $t \geq t_0 + T$, the inequality $\|\gamma_0\| \leq \delta$ implies $\|\gamma(t, t_0, \gamma_0)\| < \varepsilon$.

Definition 3.4. A function $a(r)$ is said to belong to the class K if $a \in PC([0, \psi), R_+)$, $a(0) = 0$, and $a(r)$ is strictly monotone increasing in r .

In this paper, we define the following sets:

$$\bar{S}_\psi = \{ \gamma \in R^N : \|\gamma\| \leq \psi \}$$

$$S_\psi = \{ \gamma \in R^N : \|\gamma\| < \psi \}$$

Suffice to say that the inequalities between vectors are understood to be component-wise inequalities.

We will use the comparison results for the impulsive Caputo fractional differential equation of the type



$$\begin{aligned}
 {}^C D_{t_0}^\beta u &= g(t, u), \quad t \geq t_0, \quad t \neq t_k, \quad k = 1, 2, \dots \\
 \Delta u &= \psi_k(u(t_k)), \quad t = t_k, \quad k \in N, \\
 u(t_0^+) &= u_0,
 \end{aligned} \tag{3.5}$$

existing for $t \geq t_0$, $u \in R^n$, $R_+ = [t_0, \infty)$, $g : R_+ \times R_+^n \rightarrow R^n$, $g(t, 0) \equiv 0$, where g is the continuous mapping of $R_+ \times R_+^n$ into R^n . The function $g \in PC[R_+ \times R_+^n, R^n]$ is such that for any initial data $(t_0, u_0) \in R_+ \times R^n$, the system (3.5) with initial condition $u(t_0) = u_0$ is assumed to have a solution $u(t, t_0, u_0) \in PC^\beta([t_0, \infty), R^n)$.

Lemma 3.2. Assume $m \in PC([t_0, T] \times \bar{S}_\psi, R^N)$. and suppose there exists $t^* \in [t_0, T]$ such that for $\alpha_1 < \alpha_2$, $m(t^*, \alpha_1) = m(t^*, \alpha_2)$ and $m(t, \alpha_1) < m(t, \alpha_2)$ for $t_0 \leq t < t^*$. Then if the Caputo fractional Dini derivative of m exists at t^* , then the inequality ${}^C D_+^\beta m(t^*, \alpha_1) - {}^C D_+^\beta m(t^*, \alpha_2) > 0$ holds.

Proof. Let $\Omega(t, \gamma) = m(t, \alpha_1) - m(t, \alpha_2)$. Applying (3.4), we have

$$\begin{aligned}
 {}^C D_+^\beta (m(t^*, \alpha_1) - m(t^*, \alpha_2)) &= \limsup_{h \rightarrow 0^+} \frac{1}{h^\beta} \{ [m(t^*, \alpha_1) - m(t^*, \alpha_2)] - [m(t_0, \alpha_1) - m(t_0, \alpha_2)] \\
 &\quad - \sum_{r=1}^{\lfloor \frac{t-t_0}{h} \rfloor} (-1)^{r+1} C_r [m(t^* - rh, \alpha_1) - m(t^* - rh, \alpha_2)] - [m(t_0, \alpha_1) - m(t_0, \alpha_2)] \}
 \end{aligned}$$

when $\alpha_1 = \alpha_2$ we have

$$\begin{aligned}
 {}^C D_+^\beta (m(t^*, \alpha_1) - m(t^*, \alpha_2)) &= \limsup_{h \rightarrow 0^+} \frac{1}{h^\beta} \{ -[m(t_0, \alpha_1) - m(t_0, \alpha_2)] \\
 &\quad - \sum_{r=1}^{\lfloor \frac{t-t_0}{h} \rfloor} (-1)^{r+1} C_r [m(t^* - rh, \alpha_1) - m(t^* - rh, \alpha_2)] - [m(t_0, \alpha_1) - m(t_0, \alpha_2)] \}
 \end{aligned}$$

$$\begin{aligned}
 {}^C D_+^\beta (m(t^*, \alpha_1) - m(t^*, \alpha_2)) &= - \limsup_{h \rightarrow 0^+} \frac{1}{h^\beta} [m(t_0, \alpha_1) - m(t_0, \alpha_2)] \\
 &\quad + \limsup_{h \rightarrow 0^+} \frac{1}{h^\beta} \sum_{r=1}^{\lfloor \frac{t-t_0}{h} \rfloor} (-1)^r C_r [m(t^* - rh, \alpha_1) - m(t^* - rh, \alpha_2)] \\
 &\quad - \limsup_{h \rightarrow 0^+} \frac{1}{h^\beta} \sum_{r=1}^{\lfloor \frac{t-t_0}{h} \rfloor} (-1)^r C_r [m(t_0, \alpha_1) - m(t_0, \alpha_2)]
 \end{aligned}$$

$$\begin{aligned}
 {}^C D_+^\beta (m(t^*, \alpha_1) - m(t^*, \alpha_2)) &= - \limsup_{h \rightarrow 0^+} \frac{1}{h^\beta} [m(t_0, \alpha_1) - m(t_0, \alpha_2)] \\
 &\quad - \limsup_{h \rightarrow 0^+} \frac{1}{h^\beta} \sum_{r=1}^{\lfloor \frac{t-t_0}{h} \rfloor} (-1)^r C_r [m(t_0, \alpha_1) - m(t_0, \alpha_2)]
 \end{aligned}$$



$${}^c D_+^\beta (m(t^*, \alpha_1) - m(t^*, \alpha_2)) = - \limsup_{h \rightarrow 0^+} \frac{1}{h^\beta} \sum_{r=0}^{\lfloor \frac{t-t_0}{h} \rfloor} (-1)^r \beta C_r [m(t_0, \alpha_1) - m(t_0, \alpha_2)]$$

Applying (3.8) in [1], we have

$${}^c D_+^\beta m(t^*, \alpha_1) - {}^c D_+^\beta m(t^*, \alpha_2) = - \frac{(t-t_0)^{-\beta}}{\Gamma(1-\beta)} [m(t^*, \alpha_1) - m(t^*, \alpha_2)]$$

By the lemma, we have that

$$m(t, \alpha_1) - m(t, \alpha_2) < 0 \text{ for } t_0 \leq t \leq t^*$$

And so, it follows that

$${}^c D_+^\beta m(t^*, \alpha_1) - {}^c D_+^\beta m(t^*, \alpha_2) > 0$$

Note that some existence results for (3.5) are given in [11, 13-14].

Remark 3.3. Lemma 3.2 extends Lemma 1 in [1], where the vectors $m(t, \alpha_1)$ and $m(t, \alpha_2)$ are compared component-wise.

In the following, we establish the comparison result for the system (3.3).

4. Fractional differential inequalities and comparison results

For vector fractional differential equation

In this section, again we assume that $0 < \beta < 1$.

Theorem 4.1 (Comparison Result). Assume that:

(i) $g \in PC[R_+ \times R_+^n, R^n]$ and is continuous in $(t_{k-1}, t_k]$, $k = 1, 2, \dots$ and $g(t, u)$ is quasimonotone non-decreasing in u for each $u \in R^n$ and $\lim_{(t, \gamma) \rightarrow (t_k^+, u)} g(t, u) = g(t_k^+, u)$ exists;

(ii) $\Omega \in PC[R_+ \times R^N, R^N]$ and $\Omega \in \chi$ such that

$${}^c D_+^\beta \Omega(t, \gamma) \leq g(t, \Omega(t, \gamma)), \quad (t, \gamma) \in R_+ \times R^N$$

and

$\Omega(t_k, \gamma + I_k(\gamma(t_k))) \leq \omega_k(\Omega(t, \gamma(t))), t = t_k, \gamma \in S_\psi$ and the function $\omega_k : R_+^N \rightarrow R_+^N$ is nondecreasing for $k = 1, 2, \dots$



(iii) $\mathcal{G}(t) = \mathcal{G}(t, t_0, u_0) \in PC^\beta([t_0, T], R^n)$ be the maximal solution of the IVP for the IFRDE system (3.5)

Then,

$$\Omega(t, \gamma(t)) \leq \mathcal{G}(t), \quad t \geq t_0, \quad (4.1)$$

where $\gamma(t) = \gamma(t, t_0, \gamma_0) \in PC^\beta([t_0, T], R^N)$ is any solution of (3.3) existing on $[t_0, \infty)$, provided that

$$\Omega(t_0^+, \gamma_0) \leq u_0 \quad (4.2)$$

Proof.

Let $\eta \in S_\psi$ be a small enough arbitrary vector and consider the initial value problem for the following system of fractional differential equations,

$$\begin{aligned} {}^c D^\beta u &= g(t, u) + \eta, \quad \text{for } t \in [t_0, \infty) \\ u(t_0^+) &= u_0 + \eta, \end{aligned} \quad (4.3)$$

for $t \in [t_0, \infty)$.

The function $u_\eta(t, \alpha)$ is a solution of (4.3), where $\alpha > 0$ the fractional differential equation (3.5) if and only if it satisfies the Volterra fractional integral equation,

$$u_\eta(t, \alpha) = u_0 + \eta + \frac{1}{\Gamma(\beta)} \int_{t_0}^t (t-s)^{\beta-1} (g(s, u_\eta(s, \alpha)) + \alpha) ds, \quad t \in [t_0, \infty). \quad (4.4)$$

Let the function $m(t, \alpha) \in PC([t_0, T] \times S_\psi, R_+^N)$ be defined as $m(t, \alpha) = \Omega(t, \gamma^*(t))$

We now prove that

$$m(t, \alpha) < u_\eta(t, \alpha) \quad \text{for } t \in [t_0, \infty) \quad (4.5)$$

Observe that the inequality (4.5) holds whenever $t = t_0$ i.e.

$$m(t_0, \alpha) = \Omega(t_0, \gamma_0) \leq u_0 < u_\eta(t_0, \alpha)$$

Assume that the inequality (4.5) is not true, then there exists a point $t_1 > t_0$ such that

$$m(t_1, \alpha) = u_\eta(t_1, \alpha) \quad \text{and} \quad m(t, \alpha) < u_\eta(t, \alpha) \quad \text{for } t \in [t_0, t_1).$$

It follows from Lemma 3.2 that

$${}^c D_+^\beta (m(t_1, \alpha) - u_\eta(t_1, \alpha)) > 0 \quad \text{i.e.}$$



$${}^C D_+^\beta (m(t_1, \alpha)) > {}^C D_+^\beta u_\eta(t_1, \alpha)$$

$${}^C D_+^\beta \Omega(t_1, \gamma(t_1)) > {}^C D_+^\beta u_\eta(t_1, \alpha)$$

and using (4.3) we arrive at

$${}^C D_+^\beta \Omega(t_1, \gamma(t_1)) > g(t_1, u(t_1, \alpha)) + \eta > g(t_1, u(t_1, \alpha))$$

Therefore,

$${}^C D_+^\beta m(t_1, \alpha) > g(t_1, u(t_1, \alpha)) \quad (4.6)$$

From Theorem 4.1, the function $\gamma^*(t) = \gamma(t, t_0, \gamma_0)$ satisfies the IVP (4.3) and the equality

$$\limsup_{h \rightarrow 0^+} \frac{1}{h^\beta} [\gamma^*(t) - \gamma_0 - S(\gamma^*(t), h)] = f(t, \gamma^*(t)), \text{ holds} \quad (4.7)$$

where $\gamma^*(t) = \gamma(t, t_0, \gamma_0)$ is any other solution of (3.5), and

$$S(\gamma^*(t), h) = \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} C_r [\gamma^*(t - rh) - \gamma_0] \quad (4.8)$$

is the Grunwald Letnikov fractional derivative

Multiply equation (4.7) through by h^β

$$\limsup_{h \rightarrow 0^+} [\gamma^*(t) - \gamma_0 - S(\gamma^*(t), h)] = h^\beta f(t, \gamma^*(t))$$

$$\gamma^*(t) - \gamma_0 - [S(\gamma^*(t), h) + \rho(h^\beta)] = h^\beta f(t, \gamma^*(t))$$

$$\gamma^*(t) - h^\beta f(t, \gamma^*(t)) = [S(\gamma^*(t), h) + \gamma_0 + \rho(h^\beta)] \quad (4.9)$$

Then for $t \in [t_0, \infty)$, we have



$$\begin{aligned}
 & m(t, \alpha) - m(t_0, \alpha) - \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} \beta C_r [m(t-rh, \alpha) - m(t_0, \alpha)] = \\
 & \Omega(t, \gamma^*(t)) - \Omega(t_0, \gamma_0) - \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} \beta C_r [\Omega(t-rh, \gamma^*(t)) - h^\beta f(t, \gamma^*(t)) - \Omega(t_0, \gamma_0)] \\
 & = \Omega(t, \gamma^*(t)) - \Omega(t_0, \gamma_0) - \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} \beta C_r [\Omega(t-rh) - \gamma^*(t) - h^\beta f(t, \gamma^*(t)) - \Omega(t_0, \gamma_0)] + \\
 & \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} \beta C_r \{ [\Omega(t-rh), S(\gamma^*(t), h) + \gamma_0 + \rho(h^\beta) - \Omega(t_0, \gamma_0)] - [\Omega(t-rh, \gamma^*(t-rh)) - \Omega(t_0, \gamma_0)] \}
 \end{aligned}$$

(4.10)

Since $\Omega(t, \gamma)$ is locally Lipschitzian with respect to the second variable, we have that,

$$\begin{aligned}
 & \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} \beta C_r \{ [\Omega(t-rh), S(\gamma^*(t), h) + \gamma_0 + \rho(h^\beta)] - [\Omega(t-rh, \gamma^*(t-rh))] \} \\
 & \leq L \left| (-1)^{r+1} \left\| \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (\beta C_r) [S(\gamma^*(t), h) + \gamma_0 + \rho(h^\beta) - \gamma^*(t-rh)] \right\| \right|,
 \end{aligned}$$

where $L > 0$ is a Lipschitz constant

$$\leq L \left\| \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (\beta C_r) [S(\gamma^*(t), h) + \rho(h^\beta)] - (\gamma^*(t-rh) - \gamma_0) \right\|. \tag{4.11}$$

Using (4.8), equation (4.11) becomes,

$$\begin{aligned}
 & \leq L \left\| \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (\beta C_r) \left(\sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} (\beta C_r) [(\gamma^*(t-rh) - \gamma_0) + \rho(h^\beta)] - (\gamma^*(t-rh) - \gamma_0) \right) \right\| \\
 & \leq L \left\| \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (\beta C_r) (-1)^{r+1} \left[\sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} \beta C_r [(\gamma^*(t-rh) - \gamma_0)] + \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} \beta C_r \rho(h^\beta) - \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} \beta C_r (\gamma^*(t-rh) - \gamma_0) \right] \right\| \\
 & \leq L (-1)^{r+1} \left\| \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (\beta C_r) [(\gamma^*(t-rh) - \gamma_0)] + \left[\sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} \beta C_r - 1 \right] + \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} \beta C_r \rho(h^\beta) \right\|. \tag{4.12}
 \end{aligned}$$

Substituting (4.12) into (4.10) we have



$$\begin{aligned}
 &= \Omega(t, \gamma^*(t)) - \Omega(t_0, \gamma_0) - \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} \beta C_r \left[\Omega(t-rh) - \gamma^*(t) - h^\beta f(t, \gamma^*(t)) - \Omega(t_0, \gamma_0) \right] + \\
 &L \left\| \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} (\beta C_r) (\gamma^*(t-rh) - \gamma_0) \left[\sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} \beta C_r - 1 \right] + \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} \beta C_r \rho(h^\beta) \right\| \\
 &= \Omega(t, \gamma^*(t)) - \Omega(t_0, \gamma_0) - \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} \beta C_r \left[\Omega(t-rh) - \gamma^*(t) - h^\beta f(t, \gamma^*(t)) - \Omega(t_0, \gamma_0) \right] + \\
 &L \left\| \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} (\beta C_r) (\gamma^*(t-rh) - \gamma_0) \left[- \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^r \beta C_r - 1 \right] + \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} \beta C_r \rho(h^\beta) \right\|
 \end{aligned}$$

Dividing through by $h^\beta > 0$ and taking the \limsup as $h \rightarrow 0^+$ we have,

$$\begin{aligned}
 {}^c D_+^\beta m(t, \alpha) &= \limsup_{h \rightarrow 0^+} \frac{1}{h^\beta} \left\{ \Omega(t, \gamma^*(t)) - \Omega(t_0, \gamma_0) - \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} \beta C_r \left[\Omega(t-rh) - \gamma^*(t) - h^\beta f(t, \gamma^*(t)) - \Omega(t_0, \gamma_0) \right] \right\} + \\
 &\limsup_{h \rightarrow 0^+} \frac{1}{h^\beta} L \left\| \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} (\beta C_r) (\gamma^*(t-rh) - \gamma_0) \left[- \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^r \beta C_r - 1 \right] + \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} \beta C_r \rho(h^\beta) \right\|.
 \end{aligned}$$

Recall that,

$$\lim_{h \rightarrow 0^+} \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} (\beta C_r) = -1 \text{ and } \limsup_{h \rightarrow 0^+} \frac{1}{h^\beta} \rho(h^\beta) = 0.$$

From (3.6) and (3.7) in [1], we have that

$${}^c D_+^\beta m(t, \alpha) = {}^c D_+^\beta \Omega(t, \gamma^*(t)) + L \left\| \sum_{r=1}^{\lceil \frac{t-t_0}{h} \rceil} (-1)^{r+1} (\beta C_r) (\gamma^*(t-rh) - \gamma_0) [-(-1) - 1] + 0 \right\|.$$

Using condition (ii) of the Theorem 4.1, we obtain the estimate

$${}^c D_+^\beta m(t, \alpha) \leq g(t, \Omega(t, \gamma^*(t))) = g(t, m(t, \alpha)), \tag{4.13}$$

Also,

$$m(t_0^+, \alpha) \leq u_0 \text{ and } m(t_k^+, \alpha) = \Omega(t_k^+, \gamma(t_k)) + I_k(\gamma(t_k)) \leq \rho_k(m(t_k)) \tag{4.14}$$

Now (4.14) with $t = t_1$ contradicts (4.6), hence (4.5) is true. □

For $t \in [t_0, T]$, we now establish that



$$u_{\eta_1}(t, \alpha) < u_{\eta_2}(t, \alpha) \quad \text{whenever } \eta_1 < \eta_2 \tag{4.15}$$

Observe that the inequality (4.15) holds for $t = t_0$

Assume that (4.15) is not true. Then there exists a point t_1 such that $u_{\eta_1}(t_1, \alpha) = u_{\eta_2}(t_2, \alpha)$ and $u_{\eta_1}(t, \alpha) < u_{\eta_2}(t, \alpha)$ for $t \in [t_0, t_1)$.

By Lemma 3.2, we have that

$${}^c D_+^\beta (u_{\eta_1}(t_1, \alpha) - u_{\eta_2}(t_2, \alpha)) > 0$$

However,

$$\begin{aligned} {}^c D_+^\beta (u_{\eta_1}(t_1, \alpha) - u_{\eta_2}(t_2, \alpha)) &= {}^c D_+^\beta (u_{\eta_1}(t_1, \alpha) - u_{\eta_2}(t_2, \alpha)) \\ &= g(t_1, u(t_1, \alpha)) + \eta_1 - [g(t_1, u(t_1, \alpha)) + \eta_2] \\ &= \eta_1 - \eta_2 < \mathbf{0} \end{aligned}$$

which is a contradiction, and so (4.15) is true. Thus, equations (4.5) and (4.15) guarantee that the family of solutions $\{u_\eta(t, \alpha)\}$, $t \in [t_0, T]$ of (4.3) is uniformly bounded, i.e. there exists $\lambda > 0$ with $|u_\eta(t, \alpha)| \leq \lambda$, with bound λ on $[t_0, T]$. We now show that the family $\{u_\eta(t, \alpha)\}$ is equicontinuous on $[t_0, T]$. Let $k = \sup\{|g(t, \gamma)| : (t, \gamma) \in [t_0, T] \times [-\lambda, \lambda]\}$, where λ is the bound on the family $\{u_\eta(t, \alpha)\}$. Fix a decreasing sequence $\{\eta_i\}_{i=0}^\infty(t)$, such that $\lim_{i \rightarrow \infty} \eta_i = 0$ and consider a sequence of functions $\{u_\eta(t, \alpha)\}$. Again, let $t_1, t_2 \in [t_0, T]$, with $t_1 < t_2$, then we have the following estimates,

$$\begin{aligned} \|u_\eta(t_2, \alpha) - u_\eta(t_1, \alpha)\| &\leq \|u_0 + \eta_i + \frac{1}{\Gamma(\beta)} \int_{t_0}^{t_2} (t_2 - s)^{\beta-1} (g(s, u(s, \eta_i)) + \eta_i) ds \\ &\quad - \left(u_0 + \eta_i + \frac{1}{\Gamma(\beta)} \int_{t_0}^{t_1} (t_1 - s)^{\beta-1} (g(s, u(s, \eta_i)) + \eta_i) ds \right) \| \\ &\leq \frac{1}{\Gamma(\beta)} \left\| \left[\int_{t_0}^{t_2} (t_2 - s)^{\beta-1} - \int_{t_0}^{t_1} (t_1 - s)^{\beta-1} \right] (g(s, u(s, \eta_i)) + \eta_i) ds \right\| \\ &\leq \frac{k}{\Gamma(\beta)} \left\| \left(\int_{t_0}^{t_2} (t_2 - s)^{\beta-1} - \int_{t_0}^{t_1} (t_1 - s)^{\beta-1} \right) ds \right\| \end{aligned}$$



$$\begin{aligned}
 &\leq \frac{k}{\Gamma(\beta)} \left\| \left(\int_{t_0}^{t_1} (t_2 - s)^{\beta-1} + \int_{t_1}^{t_2} (t_2 - s)^{\beta-1} - \int_{t_0}^{t_1} (t_1 - s)^{\beta-1} \right) ds \right\| \\
 &\leq \frac{k}{\Gamma(\beta)} \left\| \int_{t_0}^{t_1} (t_2 - s)^{\beta-1} - \int_{t_0}^{t_1} (t_1 - s)^{\beta-1} + \int_{t_1}^{t_2} (t_2 - s)^{\beta-1} ds \right\| \\
 &\leq \frac{k}{\Gamma(\beta)} \left[\left\| \int_{t_0}^{t_1} (t_2 - s)^{\beta-1} - \int_{t_0}^{t_1} (t_1 - s)^{\beta-1} ds \right\| + \left\| \int_{t_1}^{t_2} (t_2 - s)^{\beta-1} ds \right\| \right] \\
 &\leq \frac{k}{\beta\Gamma(\beta)} \left[\|(t_2 - t_0)^\beta - (t_1 - t_0)^\beta - (t_2 - t_1)^\beta\| + \|(t_2 - t_1)^\beta\| \right] \\
 &\leq \frac{k}{\beta\Gamma(\beta)} \left[(t_2 - t_1)^\beta + (t_2 - t_1)^\beta \right] = \frac{2k}{\Gamma(\beta+1)} (t_2 - t_1)^\beta < \varepsilon
 \end{aligned}$$

provided $\|t_2 - t_1\| < \delta(\varepsilon) = \left(\frac{\Gamma(\beta+1)\varepsilon}{2k} \right)^{\frac{1}{\beta}}$, proving that the family of solutions $\{u_{\eta_j}(t, \alpha)\}$ is equicontinuous. By the Arzela-Ascoli theorem, $\{u_{\eta_j}(t, \alpha)\}$ guarantees the existence of a subsequence $\{u_{\eta_{i_j}}(t, \alpha)\}$ which converges uniformly to the function $\mathcal{G}(t)$ on $[t_0, T]$. Then we show that $\mathcal{G}(t)$ is a solution of (4.4). Thus, equation (4.4) becomes

$$u_{\eta_{i_j}}(t, \alpha) = u_{0_{i_j}} + \eta_{i_j} + \frac{1}{\Gamma(\beta)} \int_{t_0}^t (t-s)^{\beta-1} (g_{i_j}(s, u_{\eta_{i_j}}(s, \alpha)) + \eta_{i_j}) ds, \quad t \in [t_0, \infty). \tag{4.16}$$

Taking the lim as $i_j \rightarrow \infty$ in (4.16) yields,

$$\mathcal{G}(t) = u_0 + \eta_{i_j} + \frac{1}{\Gamma(\beta)} \int_{t_0}^t (t-s)^{\beta-1} (g(s, \mathcal{G}(t))) ds, \quad t \in [t_0, \infty). \tag{4.17}$$

Hence, $\mathcal{G}(t)$ is a solution of (3.5) on $[t_0, T]$. We claim that $\mathcal{G}(t)$ is the maximal solution of (3.5). Then, from (4.5), we have that $m(t, \alpha) < u_{\eta_j}(t, \alpha) \leq \mathcal{G}(t)$ on $[t_0, T]$.

Suppose that in Theorem 4.1, $g(t, u) \equiv 0$, then we have the following results

Corollary 4.1.

Assume that Condition (i) of Theorem 4.1 holds and,

- (i) $\Omega \in PC[R_+ \times R^N, R_+^N]$ and $\Omega \in \chi$ such that

$${}^c D_+^\beta \Omega(t, \gamma) \leq 0$$

(4.14) holds, and



$\Omega(t_k, \gamma + I_k(\gamma(t_k))) \leq \omega_k(\Omega(t, \gamma(t))), t = t_k, \gamma \in S_\psi$ and the function $\omega_k : R_+^N \rightarrow R_+^N$ is nondecreasing for $k = 1, 2, \dots$

Then for $t \in [t_0, \infty)$, the inequality

$$\Omega(t, \gamma(t)) \leq \Omega(t_0^+, \gamma_0) \text{ holds}$$

MAIN RESULTS

In this section, we will obtain sufficient conditions for the practical stability of the system (3.3). Again we assume $0 < \beta < 1$.

Theorem 5.1. Assume that:

(i) $g \in PC[R_+ \times R_+^n, R^n]$ is piecewise continuous in $(t_{k-1}, t_k]$ and for each $u \in R^n, k = 1, 2, \dots$, and $\lim_{(t,y) \rightarrow (t_k^+, u)} g(t, y) = g(t_k^+, u)$ exists, and $g(t, u)$ is quasimonotone nondecreasing in u

(ii) $\Omega \in PC[R_+ \times R^N, R_+^N]$ and $\Omega \in \chi$ such that

$${}^c D_+^\beta \Omega(t, \gamma) \leq g(t, \Omega(t, \gamma)), t \neq t_k, \text{ holds for all } (t, \gamma) \in R_+ \times S_\psi,$$

There exists $\psi_0 > 0$ such that $\gamma_0 \in S_\psi$ implies that $\gamma(t) + I_k(\gamma(t_k)) \in S_\psi$ and

$\Omega(t_k, \gamma + I_k(\gamma(t_k))) \leq \omega_k(\Omega(t, \gamma(t))), t = t_k, \gamma \in S_\psi$ and the function $\omega_k : R_+^N \rightarrow R_+^N$ is nondecreasing for $k = 1, 2, \dots$

(iii) $b(\|\gamma\|) \leq \Omega_0(t, \gamma)$, where $b \in K$ and $\Omega_0(t, \gamma) = \sum_{i=1}^N \Omega_i(t, \gamma)$.

Then the practical stability of the trivial solution $u = 0$ of (3.5) implies the practical stability of the trivial solution $\gamma = 0$ of (3.3).

Proof. Let $0 < \varepsilon < \rho$ and $t_0 \in R_+$ be given.

Assume that the solution (3.5) is stable. Then given $b(\varepsilon) > 0$ and $t_0 \in R_+$, there exists a positive function $\delta = \delta(t_0, \varepsilon) > 0$ which is continuous in t_0 for each ε such that

$$\sum_{i=1}^N u_{i0} < \delta \text{ implies } \sum_{i=1}^N u_i(t, t_0, u_0) < b(\varepsilon), t \geq t_0 \tag{5.1}$$

where $u(t, t_0, u_0)$ is any solution of (3.5).



Choose $u_0 = \Omega(t_0^+, \gamma_0)$.

Since $\Omega(t, x)$ is continuous, then by the property of continuity, given $\varepsilon > 0$ there exists a positive function $\delta_1 = \delta_1(t_0, \delta) > 0$ that is continuous in t_0 for each δ such that the inequalities

$$\|\Omega(t, \gamma) - \Omega(t_0, \gamma_0)\| < \delta \text{ implies } \|\gamma - \gamma_0\| < \delta_1$$

and as $\|\Omega(t, \gamma)\| \rightarrow 0, \|\gamma\| \rightarrow 0$ then the inequalities

$$\|\gamma_0\| < \delta_1 \text{ and } \|\Omega(t_0, \gamma_0)\| < \delta \tag{5.2}$$

are satisfied simultaneously.

We claim that, if $\|\gamma_0\| < \delta_1$ then $\|\gamma(t, t_0, \gamma_0)\| < \varepsilon$.

Suppose that this claim is false, then there would exist a point $t_1 \in [t_0, t)$ and the solution $\gamma(t, t_0, \gamma_0)$ with $\|\gamma_0\| < \delta_1$ such that

$$\|\gamma(t_1)\| = \varepsilon \text{ and } \|\gamma(t)\| < \varepsilon \text{ for } t \in [t_0, t_1) \tag{5.3}$$

So that using equation (5.3); condition (iii) of Theorem 5.1 reduces to the form

$$b(\|\gamma(t_1)\|) \leq \sum_{i=1}^N \Omega_i(t_1, \gamma(t_1)), \text{ implying}$$

$$b(\varepsilon) \leq \sum_{i=1}^N \Omega_i(t_1, \gamma(t_1)) \tag{5.4}$$

for $t \in [t_0, t_1)$ and from Theorem 4.1,

$$\Omega(t, \gamma(t)) \leq \mathcal{G}(t) \tag{5.5}$$

where $\mathcal{G}(t) = \sum_{i=1}^n \mathcal{G}_i(t, t_0, u_0)$ is the maximal solution of (3.5).

Then, using equations (5.4), (5.3), (5.5) and condition (iii) of Theorem 5.1 we arrive at the estimate

$$b(\varepsilon) \leq \Omega_0(t_1, \gamma(t_1)) \leq \sum_{i=1}^N \mathcal{G}_i(t, t_0, u_0) < b(\varepsilon)$$

which leads to a contradiction.

Hence, the practical stability of the trivial solution $u = 0$ of (3.5) implies the practical stability of the trivial solution $\gamma = 0$ of (3.3).



CONCLUSION

In this paper, the existence of the maximal solution of the comparison system for vector Lyapunov function is established, and sufficient conditions for the practical stability of impulsive Caputo fractional order systems are presented by means of the comparison principle. It was discovered that the vector form of the Lyapunov function used was majorized by the maximal solution of the comparison system. The sufficient conditions for the practical stability of the comparison impulsive fractional order system as established went further to illustrate the fact that the solution of the main system is practically stable in the sense of Lyapunov.

REFERENCES

- [1] R. Agarwal, D. O'Regan, and S. Hristova, "Stability of Caputo Fractional Differential Equations". Applications of Mathematics. 60, No. 6, 653-676, 2015.
- [2] R. Agarwal, S. Hristova and D. O'Regan, "Stability of Solutions of Impulsive Caputo Fractional Differential Equations," Electronic Journal of Differential Equations, vol. 2016, No. 58, pp. 1-22, 2016.
- [3] R. Agarwal, D. O'Regan, S. Hristova, and Cicek, M., (2017). Practical Stability with respect to Initial Time Difference for Caputo Fractional Differential Equations. Electronic Journal of Differential Equations. Communications in Nonlinear Science and Numerical Simulations, 42, 106, 2017, doi.org/10.1016/j.cnsns.2016.05.005.
- [4] E. P. Akpan and O. Akinyele, "On the ϕ_0 -Stability of Comparison Differential Systems." Journal of Mathematical Analysis and Applications, 164(2), 307-324, 1992.
- [5] L. Arnold and B. Schmalfuss, "Lyapunov's Second Method for Random Dynamical Systems." Journal of Differential Equations, 177(1), 235-265.
- [6] D. Baleanu, and O. G. Mustafa, "On the Global Existence of Solutions to a Class of Fractional Differential Equations." Comput. Math. Appl. 59. 1835-1841, 2010.
- [7] D. D. Bainov and P. S. Simeonov, "Systems with Impulse Effects: Stability, Theory and Applications," Ellis Horwood, Chichester, 1989.
- [8] D. D. Bainov and I. Stamova, "On the Practical Stability of Differential-Difference Equations with Variable Impulsive Perturbations." Journal of *Mathematical Analysis and Applications*, 200, 272-288, 1996.
- [9] T. A. Burton, "Fractional Differential Equations and Lyapunov functionals. "Nonlinear Analysis: Theory, Methods and Applications, 74(16), 5648-5662, 2011.
- [10] K. Diethelm, "The Analysis of Fractional Differential Equations. An Application Oriented Exposition Using Differential Operators of Caputo type." Springer, Berlin, (2010).
- [11] J. V. Devi, F. A. Mc Rae, and Z. Drici, "Variational Lyapunov method for Fractional Differential Equations," *Comput. Math. Appl.*, 64, 2982-2989, 2012.
- [12] D. K. Igobi and J. E. Ante, "Results on Existence and Uniqueness of Solutions of Impulsive Neutral Integro-Differential System". Journal of Mathematics Research, Vol 10, No. 4, 2018, <http://doi:10.5539/jmr.v10n4p165>.
- [13] D. K. Igobi and M. P. Ineh, "Results on Existence and Uniqueness of Solutions of Dynamic Equation on Time Scale via Generalized Ordinary Differential Equations." International Journal of Applied Mathematics, 37, No. 1, 1-20, 2024.



- [14] A. A. Kilbas, Marichev, O. I. and S. G. Samko, "Fractional Integrals and Derivatives (Theory and Applications), Transl. from the Russian. Gordon and Breach, New York, 1974.
- [15] V. Lakshmikantham, "On the Method of vector Lyapunov Functions," 1974, Technical report No. 16.
- [16] V. Lakshmikantham, D. D. Bainov and P. S. Simeonov, "The Theory of Impulsive Differential Equations. Series in Modern Applied Mathematics, No. 6, 1989, World Scientific Publishing C. Pte, Ltd (2009).
- [17] V. Lakshmikantham and A. S. Vatsala, "Basic Theory of Impulsive Differential Equations," *Nonlinear Analysis: Theory, Methods & Applications*, 69(8), 2677-2682, 2008.
- [18] V. Lakshmikantham, S. Leela and M. Sambandham, "Lyapunov Theory for Fractional Differential Equations," *Communications in Applied Analysis*, 12 (4), 365, 2008.
- [19] V. Lakshmikantham and S. Leela, "Differential and Integral Inequalities", Theory and Application, Vol. 1. Academic Press, Inc. New York and London, 1969.
- [20] V. Lakshmikantham, S. Leela and A. A. Martynyuk, "Practical Stability of nonlinear Systems, World Scientific, Singapore, 1960.
- [21] V. Lakshmikantham, S. Leela and J. V. Devi, "Theory of Fractional Dynamic Systems. Cambridge Scientific Publishers, 2009.
- [22] V. Lakshmikantham and A. S. Vatsala, "General Uniqueness and Monotone Iterative Technique for Fractional Differential Equations," *Applied Mathematics Letters* 21, 828-834, 2007.
- [23] V. Lakshmikantham, V. M. Matrosov and S. Sivasundaram, "On Vector Lyapunov Functions and Stability Analysis of Nonlinear System," Springer Science-Business Media Dordrecht, 1991.
- [24] J. Lasalle and S. Lefschetz, "Stability by Liapunov's Direct Method with Applications," Academic Press, New York, NY, USA, 1961.
- [25] Y. Li, Y. Chen, I. Podlubny, "Stability of fractional order nonlinear dynamic systems: Lyapunov Direct Method and Generalized Mittag-leffler Stability." *Comput. Math. Appl.* 59, 1810-1821, 2010.
- [26] C. Li, D. Qian, Y. Chen, "On Riemann-Liouville and Caputo derivatives." *Discrete Dyn. Nat. Soc.* Article ID 562494, 15 pages, 2011.
- [27] C. P. Li and F. R. Zhang, "A survey on the Stability of Fractional Differential Equations," *Eur. Phys. J. Special Topics* 193, 27-47, 2011, doi:10.1140/epjst/e2011-01379-1.
- [28] Z. Lipcsey, J. A. Ugboh, I. M. Esuabana and I. O. Isaac, "Existence Theorem for Impulsive Differential Equations with Measurable Right Side for Handling Delay Problems." *Hindawi Journal of Mathematics*, Vol. 2020, Article ID 7089313, 17 pages, 2020
- [29] A. Ludwig, B. Pustal, D. M. Herlach, "General Concept for a Stability Analysis of a Planar Interface under Rapid Solidification Conditions in Multi-Component Alloy Systems." *Material Science and Engineering: A*, 304, 277-280, 2001.
- [30] V. D. Milman and A. D. Myshkis, "On Motion Stability with Shocks," *Sibirsk. Mat Zh.* 1 233-237, 1960, (in Russian).
- [31] A. A. Martynyuk, "Practical Stability of Motion," Nauka Dumka Kiev, 1983 [Russian].
- [32] A. A. Martynyuk, "Practical Stability Conditions for Hybrid Systems," In 12th World Congress of IMACS, Paris, pp. 344-347, 1988.



- [33] A. A. Martynyuk, "On Practical Stability of Hybrid Systems," *Appl. Mech.* 25, 101-107, 1989 [Russian].
- [34] I. Podlubny, "Fractional differential equations. An Introduction to Fractional Derivatives, Fractional Differential Equations to methods of their Solutions and some of their Applications. *Mathematics in Science and Engineering 198*, Academic Press, San Diego, 1999.
- [35] Z. Qunli, "A Class of Vector Lyapunov Functions for Stability Analysis of Nonlinear Impulsive Differential System," Hindawi Publishing Corporation. *Mathematical problems in Engineering*, Article ID 649012, 9 pages, 2014.
- [36] B. S. Razumikhin, "Stability of Systems with Retardation, Nauka, Moscow, [Russian], 1988.
- [37] S. G. Samko, A. A. Kilbas and O. I. Marichev, "Fractional Integrals and Derivatives : Theory and Applications," (Gordon and Breach, Amsterdam)1993.
- [38] S. K. Srivastava, K. Amanpreet, "A New Approach to Stability of Impulsive Differential Equations. *Int. Journal of Math. Analysis*, Vol. 3, no. 4, 179-185, 2009.
- [39] J. A. Ugboh and I. M. Esuabana, "Existence and Uniqueness Result for a Class of Impulsive Delay Differential Equations." *International Journal of Chemistry, Mathematics and Physics*, Vol. 2, Issue 4, 2018.
- [40] B. Wu, J. Han and X. Cai, "On the Practical Stability of Impulsive Differential Equations with Infinite Delay in Terms of Two Measures." *Abstract and Applied Analysis*, Vol. 2012, 8 pages, 2012, doi:10.1155/2012/434137.
- [41] C. Wu, "A General Comparison Principle for Caputo Fractional-Order Ordinary Differential Equations," *Fractals*, 28(04), 2050070.